

Spherical continuous wavelet transforms arising from sections of the Lorentz group

Milton Ferreira

Department of Mathematics,
University of Aveiro,
3810-193 Aveiro, Portugal.
mferreira@mat.ua.pt

Abstract

We consider the conformal group of the unit sphere S^{n-1} , the so-called proper Lorentz group $\text{Spin}^+(1, n)$, for the study of spherical continuous wavelet transforms (CWT). Our approach is based on the method for construction of general coherent states associated to square integrable group representations over homogeneous spaces. The underlying homogeneous space is an extension to the whole of the group $\text{Spin}^+(1, n)$ of the factorization of the gyrogroup of the unit ball by an appropriate gyro-subgroup. Sections on this homogeneous space are constituted by rotations of the subgroup $\text{Spin}(n)$ and Möbius transformations of the type $\varphi_a(x) = (x - a)(1 + ax)^{-1}$, where a belongs to a given section on a homogeneous space of the unit ball. This extends in a natural way the work of Antoine and Vandergheynst to anisotropic conformal dilations.

MSC 2000: Primary: 42C40; Secondary 20N05, 30G35.

Keywords: Spherical continuous wavelet transforms, Möbius transformations, gyrogroups, homogeneous spaces, sections, anisotropic dilations.

1 Introduction

In the last decade there has been a growing interest in wavelet theory on the unit sphere for an efficient analysis of spherical data. Several wavelet transforms on the sphere have been proposed in recent years and their performance depends on the application. In 1995, Schröder and Sweldens [28] describe an orthogonal wavelet transform on the sphere based on the Haar wavelet function but this construction is limited to the Haar function. Holschneider [24] was the first to build a genuine spherical CWT, but it contains a parameter that although interpreted as a dilation parameter it is difficult to compute. A satisfactory approach was obtained by Antoine and Vandergheynst [8, 9] building a rigorous spherical CWT based on a group theoretical approach. Using appropriate combinations of spherical harmonics, Freeden et al. [18] defined approximation kernels and introduced a discrete wavelet transform through a kind of multiresolution on S^2 on the Fourier domain. As demonstrated in Antoine et al. [7], the approximation scheme of Freeden et al. [18] can

be realized in the general continuous wavelet theory, based on group-theoretical considerations. Many other papers [11, 21] describe applications of spherical wavelets for detection of non-Gaussianity in Cosmic Microwave Background (CMB). These works have been extended to directional wavelet transforms [7, 22, 23, 25]. All these new continuous wavelet decompositions are interesting for data analysis. Recently, new multiscale transforms were introduced on the sphere based on curvelets and ridgelets [1, 2] with applications to denoising and filtering methods. It is visible a growing interest in directional or anisotropic wavelets on the sphere. The anisotropic dilation operator introduced in Hobson et al. [22] and used in Frossard et al. [20] allows the construction of anisotropic wavelets through dilations in two orthogonal directions. However, in this anisotropic setting, a bounded admissibility integral cannot be determined (even in the plane) and thus the synthesis of a signal from its coefficients cannot be performed. This is a direct consequence of the absence of a group structure and an easy evaluation of the proper measure in the reconstruction formula. However, in our approach we are able to derive anisotropic conformal dilations from Möbius transformations, which are natural transformations on the unit sphere and, hence, to construct anisotropic spherical CWT using group theory. Our anisotropic dilation operator is clearly different of the one defined in [22] since it depends on the section chosen on an appropriate homogeneous space of the unit ball. Thus, a whole family of anisotropic dilation operators can be defined and used in signal processing on the unit sphere.

The construction of integral transforms related to group representations on L^2 -spaces on manifolds is a non-trivial problem because the square integrability property may fail to hold. For example, in the case of the unit sphere S^{n-1} there are no square integrable unitary representations on $L^2(S^{n-1})$ for $n > 3$, [29]. A way to overcome this fact is to make the group smaller, i.e., to factor out a suitable closed subgroup H and to restrict the representation to a quotient $X = G/H$. This approach was successfully realized in the case of the unit sphere [8, 9]. The authors used the conformal group of S^{n-1} , the connected component of the Lorentz group $SO_0(1, n)$ and its Iwasawa decomposition, or KAN -decomposition, where K is the maximal compact subgroup $SO(n)$, $A = SO(1, 1) \cong (\mathbb{R}_*^+, \times)$ is the subgroup of Lorentz boosts in the x_n -direction and $N \cong \mathbb{R}^{n-1}$ is a nilpotent subgroup. The homogeneous space $X = SO_0(1, n)/N \cong SO(n) \times \mathbb{R}_*^+$ gives rise to the parameter space of the isotropic spherical CWT. This approach is restricted to pure dilations on the unit sphere, according to the inverse stereographic projection of dilations from the plane to the sphere [8, 12]. However, as we will show in this paper, the spherical CWT can be extended to anisotropic conformal dilations, while preserving the main properties of square-integrability and invertibility. Our approach allows us to connect the geometry of conformal transformations on the sphere with wavelet theory.

In our paper [12] we used rotations of the group $\text{Spin}(n)$ and Möbius transformations of the form $\varphi_a(x) = (x - a)(1 + ax)^{-1}$, $a \in B^n$, being B^n the open unit ball in $\mathbb{R}^n$, as the basic and natural transformations for building a CWT on the sphere. In this paper we will make a rigorous construction of spherical continuous wavelet transforms on the unit sphere arising from sections of the Lorentz group $\text{Spin}^+(1, n)$. In physics, the Lorentz group is the classical setting for all (nongravittional) physical phenomena. The mathematical form of the kinematical laws of special relativity, Maxwell's field equations in the theory

of electromagnetics and Dirac's equation in the theory of the electron are each invariant under Lorentz transformations. In pure mathematics, the restricted (or proper) Lorentz group arises as the Möbius group, which is the symmetry group of conformal geometry on the Riemann sphere. Clifford algebra provides an easy identification of the proper Lorentz group with the group $\text{Spin}^+(1, n)$, which is a double covering group of $SO_0(1, n)$ [3, 17, 13]. It can be decomposed by the Cartan decomposition KAK , where $K = \text{Spin}(n)$ and $AK \cong B^n$. Thus, the unit ball B^n encodes all the necessary information for the study of dilations on the unit sphere, through Möbius transformations of type $\varphi_a(x)$. This rests on the study of the properties of spherical caps under the action of Möbius transformations [12], which is in the spirit of the Erlangen program of F. Klein, the study of the invariants of a given geometry. It is also well-known that the group of projective transformations of the unit ball describes relativistic dynamics of some physical phenomena [19].

The paper is organized as follows. In Section 2 we present the notions of Clifford algebras needed to the readability of the paper. More details on this topic can be found e.g. in Delanghe et al. [16] and Cnops [13]. In Section 3 we describe the proper Lorentz group $\text{Spin}^+(1, n)$ in the framework of Clifford algebra and we introduce the structure of gyrogroup necessary to the description of the gyrogroup of the unit ball $(B^n, \oplus)$. In Section 4 we factorize the gyrogroup $(B^n, \oplus)$ by the gyro-subgroup $(D_{e_n}^{n-1}, \oplus)$ and we construct sections for the respective homogeneous space which are extended to the whole of the group $\text{Spin}^+(1, n)$. In Section 5 we derive the generalized (anisotropic) spherical CWT and in Section 6 we discuss its anisotropy and covariance properties.

2 Preliminaries

Let $e_1, \dots, e_n$ be an orthonormal basis of $\mathbb{R}^n$. The universal real Clifford algebra $Cl_{0,n}$ is the free algebra over $\mathbb{R}^n$ generated modulo the relation $x^2 = -|x|^2$, for $x \in \mathbb{R}^n$. The non-commutative multiplication in $Cl_{0,n}$ is governed by the rules

$$e_i e_j + e_j e_i = -2\delta_{ij}, \quad \forall i, j \in \{1, \dots, n\}.$$

For a set $A = \{i_1, \dots, i_k\} \subset \{1, \dots, n\}$, with $1 \leq i_1 < \dots < i_k \leq n$, let $e_A = e_{i_1} e_{i_2} \dots e_{i_k}$ and $e_\emptyset = 1$. Then $\{e_A : A \subset \{1, \dots, n\}\}$ is a basis for $Cl_{0,n}$. Thus, any $a \in Cl_{0,n}$ may be written as $a = \sum_A a_A e_A$, with $a_A \in \mathbb{R}$ or still as $a = \sum_{k=0}^n [a]_k$, where $[a]_k = \sum_{|A|=k} a_A e_A$ is a so-called *k-vector* ($k = 0, 1, \dots, n$). We can identify a vector $x = (x_1, \dots, x_n) \in \mathbb{R}^n$ with the one-vector $x = \sum_{j=1}^n x_j e_j$.

The product of any two vectors (or geometric product) is given by

$$xy = -\langle x, y \rangle + x \wedge y$$

where

$$\langle x, y \rangle = \sum_{j=1}^m x_j y_j = -\frac{1}{2}(xy + yx)$$

and

$$x \wedge y = \sum_{i < j} (x_i y_j - x_j y_i) e_{ij} = \frac{1}{2}(xy - yx)$$

are the scalar (or symmetric) part and the bivectorial (or antisymmetric) part of the geometric product, respectively.

There are two linear anti-automorphisms $*$ (reversion) and $-$ (conjugation) and an automorphism $'$ of $Cl_{0,n}$ defined on its basis elements $e_A = e_{i_1} e_{i_2} \cdots e_{i_k}$, by the rules:

$$(e_A)^* = (-1)^{\frac{k(k-1)}{2}} e_A, \quad \overline{e_A} = (-1)^{\frac{k(k+1)}{2}} e_A, \quad e'_A = (-1)^k e_A.$$

Thus, we have $(ab)^* = b^* a^*$, $\overline{ab} = \overline{b} \overline{a}$ and $(ab)' = a' b'$, for any $a, b \in Cl_{0,n}$. In particular, for vectors $\overline{x} = x' = -x$ and $x^* = x$. Each non-zero vector $x \in \mathbb{R}^n$ is invertible with $x^{-1} := \frac{\overline{x}}{|x|^2}$. Finite products of invertible vectors are invertible in $Cl_{0,n}$ and form the Clifford group $\Gamma(n)$ [13]. Elements $a \in \Gamma(n)$ such that $a\overline{a} = \pm 1$ form the $\text{Pin}(n)$ group - the double cover of the group of orthogonal transformations $O(n)$, and elements $a \in \Gamma(n)$ such that $a\overline{a} = 1$ form the $\text{Spin}(n)$ group - the double cover of the group of special orthogonal transformations $SO(n)$. For each $s \in \text{Spin}(n)$, the mapping $\chi_s : \mathbb{R}^n \rightarrow \mathbb{R}^n, \chi_s(x) = sx\overline{s}$ defines a rotation. As $\ker \chi = \{-1, 1\}$ the group $\text{Spin}(n)$ is a double covering group of $SO(n)$.

In Clifford algebra the group of Möbius transformations in $\mathbb{R}^n$ can be described by 2×2 -matrices with entries in $\Gamma(n) \cup \{0\}$, forming the group $\Gamma(1, n+1)$ [3, 13].

Theorem 1 (*Vahlen [13], 1904/ Alfhors [3], 1986*) *A matrix $A = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ represents a Möbius transformation in $\mathbb{R}^n$ if and only if*

1. $a, b, c, d \in \Gamma(n) \cup \{0\}$;
2. $ab^*, cd^*, c^*a, d^*b \in \mathbb{R}^n$;
3. *the pseudodeterminant of A , $\lambda = ad^* - bc^*$, is real and non zero.*

For a matrix A fulfilling the conditions of Theorem 1 we can define the respective fractional linear transformation of $\overline{\mathbb{R}}^n$ (here $\overline{\mathbb{R}}^n$ denotes the compactification of $\mathbb{R}^n$ by the point at infinity) given by $y = (ax + b)(cx + d)^{-1}$. Thus, we have the analogy with fractional-linear transformations of the complex plane $\mathbb{C}$. Due to the non-commutative character of Clifford algebras, $xy^{-1} \neq y^{-1}x$ in general. From now on we identify $e_1, \dots, e_n$ with the canonical basis in $\mathbb{R}^n$.

3 The proper Lorentz group $\text{Spin}^+(1, n)$

The full Lorentz group $G = SO(1, n)$ consists of linear homogeneous transformations of the $(n+1)$ -dimensional space under which the quadratic form $|x|^2 = x_0^2 - |\underline{x}|^2$, is invariant, where $x = (x_0, \underline{x})$ and $\underline{x} = (x_1, \dots, x_n)$. Here we identify x_0 with the time component and the n -vector $\underline{x} = (x_1, x_2, \dots, x_n)$ with the spatial component. The group of all Lorentz transformations preserving both orientation and the direction of time is called the proper orthochronous Lorentz group and it is denoted by $SO_0(1, n)$ or sometimes by $SO^+(1, n)$. It is generated by spatial rotations of the maximal compact subgroup $K =$

$SO(n)$ and hyperbolic rotations of the subgroup $A = SO(1, 1)$, accordingly to the Cartan decomposition KAK of $SO_0(1, n)$ (see [29, 34]).

The group $SO_0(1, n)$ is connected and locally compact. The coset space $X = SO_0(1, n)/K$ is the Lobachevsky space of dimension n . It can be realized in various manners, e.g. by the upper sheet of the hyperboloid $H^+ = \{x = (x_0, \underline{x}) : x_0^2 - |\underline{x}|^2 = 1, x_0 > 0\}$ or by the unit ball $B^n = \{x \in \mathbb{R}^n : |x| < 1\}$. However, for our purpose, we shall use throughout this paper this last realization.

The double covering group of $SO_0(1, n)$ is the group $\text{Spin}^+(1, n)$. In Clifford algebra it can be described by Vahlen matrices [3, 13, 27]. These matrices can be decomposed into the maximal compact subgroup $\text{Spin}(n)$ and the set of Möbius transformations of the form $\varphi_a(x) = (x - a)(1 + ax)^{-1}$, $a \in B^n$, which map the closed unit ball $\overline{B^n}$ onto itself. These are the multi-dimensional analogous transformations of the well known Möbius transformations of the complex plane $\varphi_u(z) = \frac{z - u}{1 - \bar{u}z}$, $u \in D$, which maps the unit disc $D = \{z \in \mathbb{C} : |z| < 1\}$ onto itself.

Möbius transformations $\varphi_a(x)$ can also be written as

$$\varphi_a(x) = \frac{(1 - |a|^2)x - (1 + |x|^2 - 2\langle a, x \rangle)a}{1 - 2\langle a, x \rangle + |a|^2|x|^2}, \quad (1)$$

from which we derive its real components.

The composition of two Möbius transformations of type (1) is (up to a rotation) again a Möbius transformation of the same type:

$$(\varphi_a \circ \varphi_b)(x) = q \varphi_{(1-ab)^{-1}(a+b)}(x) \bar{q}, \quad q = \frac{1 - ab}{|1 - ab|} \in \text{Spin}(n). \quad (2)$$

Thus, we can endow the unit ball B^n with a binary operation

$$b \oplus a = (1 - ab)^{-1}(a + b) = (a + b)(1 - ba)^{-1} = \varphi_{-b}(a). \quad (3)$$

We notice that on $(B^n, \oplus)$ the neutral element is zero while each element $a \in B^n$ has an inverse given by the symmetric element $-a$. This operation is non-commutative since

$$b \oplus a = \bar{q} (a \oplus b) q, \quad q = \frac{1 - ab}{|1 - ab|} \in \text{Spin}(n), \quad (4)$$

and it is also non-associative again due to the rotation induced by q .

Proposition 1 *For all $a, b, c \in B^n$ it holds*

$$a \oplus (b \oplus c) = (a \oplus b) \oplus (qc\bar{q}), \quad \text{with } q = \frac{1 - ab}{|1 - ab|} \in \text{Spin}(n). \quad (5)$$

Proof: By the associativity of the composition of Möbius transformations we have on the one hand

$$\begin{aligned} ((\varphi_c \circ \varphi_b) \circ \varphi_a)(x) &= ((q_1 \varphi_{b \oplus c} \bar{q}_1) \circ \varphi_a)(x) \\ &= q_1 \varphi_{b \oplus c}(\varphi_a(x)) \bar{q}_1 \\ &= q_1 q_2 \varphi_{a \oplus (b \oplus c)}(x) \bar{q}_2 \bar{q}_1 \\ &= q_1 q_2 \varphi_{a \oplus (b \oplus c)}(x) \bar{q}_2 \bar{q}_1, \end{aligned}$$

with $q_1 = \frac{1-cb}{|1-cb|}$ and $q_2 = \frac{1-(b\oplus c)a}{|1-(b\oplus c)a|}$.

On the other hand we have

$$\begin{aligned} (\varphi_c \circ (\varphi_b \circ \varphi_a))(x) &= (\varphi_c \circ (q_3 \varphi_{a\oplus b} \overline{q_3}))(x) \\ &= \varphi_c(q_3 \varphi_{a\oplus b}(x) \overline{q_3}) \\ &= q_3 \varphi_{\overline{q_3} c q_3}(\varphi_{a\oplus b}(x)) \overline{q_3} \\ &= q_3 q_4 \varphi_{(a\oplus b) \oplus (\overline{q_3} c q_3)}(x) \overline{q_4} \overline{q_3}, \end{aligned}$$

with $q_3 = \frac{1-ba}{|1-ba|}$ and $q_4 = \frac{1-(\overline{q_3} c q_3)(a\oplus b)}{|1-(\overline{q_3} c q_3)(a\oplus b)|}$.

Thus, $q_1 q_2 = q_3 q_4$ and $a \oplus (b \oplus c) = (a \oplus b) \oplus (q c \overline{q})$, with $q = \frac{1-ab}{|1-ab|}$. ■

Associativity occurs only in some special cases as we can see in the next Lemma.

Lemma 1 *If $a, b, c \in B^n$ such that $a//b$ or $a \perp c$ and $b \perp c$ then the operation $\oplus$ is associative, i.e.*

$$a \oplus (b \oplus c) = (a \oplus b) \oplus c.$$

Left and right cancellation laws have different properties:

Lemma 2 *For all $a, b \in B^n$ it holds*

$$(-b) \oplus (b \oplus a) = a \tag{6}$$

$$(a \oplus b) \oplus (q(-b)\overline{q}) = a, \text{ with } q = \frac{1-ab}{|1-ab|}. \tag{7}$$

Thus, $(B^n, \oplus)$ does not have a classic group structure but a structure of a gyrogroup. This structure was introduced by A. Ungar [31, 32, 33]. Gyrogroups are special loops which share remarkable analogies with groups, thus generalizing the notion of group. The first known gyrogroup was the relativistic gyrogroup that appeared in 1988 [32]. It consists of the unit ball $B^3 = \{x \in \mathbb{R}^3 : |x| < 1\}$ with Einstein's addition of relativistically admissible velocities. The notion of gyrogroup appears by the extension of the Einstein relativistic groupoid and the Thomas precession effect (related by the rotation induced by q). The gyrolanguage was adopted by Ungar since 1991. It rests on the unification of Euclidean and hyperbolic geometry in terms of the analogies shared [31].

Definition 1 ([31]) *A groupoid $(G, \oplus)$ is a gyrogroup if its binary operation satisfies the following axioms:*

- (G1) *In G there is at least one element 0 , called a left identity satisfying $0 \oplus a = a$, for all $a \in G$.*
- (G2) *For each $a \in G$ there is an element $\ominus a \in G$, called a left inverse of a , satisfying $\ominus a \oplus a = 0$.*
- (G3) *For any $a, b, c, d \in G$ there exists a unique element $\text{gyr}[a, b]c \in G$ such that the binary operation satisfies the left gyroassociative law*

$$a \oplus (b \oplus c) = (a \oplus b) \oplus \text{gyr}[a, b]c.$$

(G4) The map $\text{gyr}[a, b] : G \rightarrow G$ given by $c \mapsto \text{gyr}[a, b]c$ is an automorphism of the groupoid $(G, \oplus)$, that is $\text{gyr}[a, b] \in \text{Aut}(G, \oplus)$.

(G5) The gyroautomorphism $\text{gyr}[a, b]$ possesses the left loop property

$$\text{gyr}[a, b] = \text{gyr}[a \oplus b, b].$$

Definition 2 A gyrogroup $(G, \oplus)$ is gyrocommutative if its binary operation satisfies $a \oplus b = \text{gyr}[a, b](b \oplus a)$, for all $a, b \in G$.

Left gyrotranslations are defined by $\tau_a^l(b) = a \oplus b$ and right gyrotranslations by $\tau_a^r(b) = b \oplus a$. In a gyrogroup they have different properties.

Proposition 2 ([31]) Let $(G, \oplus)$ be a gyrogroup and let $a, b \in G$. The unique solution of the equation $a \oplus x = b$ in G is given by $x = \ominus a \oplus b$, and the unique solution of the equation $x \oplus a = b$ in G is given by $x = b \ominus \text{gyr}[b, a]a$.

One of the most important results of this theory is that the gyro-semidirect product of a gyrogroup $(G, \oplus)$ with a gyroautomorphism group $H \subset \text{Aut}(G, \oplus)$ is a group [31]. In our case the result is stated as

Proposition 3 The gyro-semidirect product between $(B^n, \oplus)$ and $\text{Spin}(n)$ is the group of pairs (s, a) where $a \in B^n$ and $s \in \text{Spin}(n)$, with operation $\times$ given by the gyro-semidirect product

$$(s_1, a) \times (s_2, b) = (s_1 s_2 q, b \oplus (\overline{s_2} a s_2)), \quad \text{with } q = \frac{1 - \overline{s_2} a s_2 b}{|1 - \overline{s_2} a s_2 b|}. \quad (8)$$

This is a generalization of the external semidirect product of groups (see [26]). We need to establish some properties between Möbius transformations and rotations.

Lemma 3 For $s \in \text{Spin}(n)$ and $a \in B^n$ we have

$$(i) \ \varphi_a(sx\overline{s}) = s\varphi_{\overline{s}as}(x)\overline{s} \quad \text{and} \quad (ii) \ s\varphi_a(x)\overline{s} = \varphi_{sa\overline{s}}(sx\overline{s}). \quad (9)$$

These properties are easily transferred to the gyrogroup $(B^n, \oplus)$.

Corollary 1 The following equalities hold

$$(i) \ (sa\overline{s}) \oplus b = s(a \oplus (\overline{s}bs))\overline{s} \quad \text{and} \quad (ii) \ s(a \oplus b)\overline{s} = (sa\overline{s}) \oplus (sb\overline{s}). \quad (10)$$

The relation $(sa\overline{s}) \oplus (sb\overline{s}) = s(a \oplus b)\overline{s}$ defines a homomorphism of $\text{Spin}(n)$ onto the gyrogroup $(B^n, \oplus)$.

From the decomposition (8) we can derive the Cartan or KAK decomposition of the group $\text{Spin}^+(1, n)$, where $K = \text{Spin}(n)$ and $A = \text{Spin}(1, 1)$ is the subgroup of Lorentz boosts in the e_n -direction.

Lemma 4 *Each $a \in B^n$ can be written as $a = sre_n \bar{s}$, where $r = |a| \in [0, 1[$ and $s = s_1 \cdots s_{n-1} \in \text{Spin}(n)$ with*

$$s_i = \cos \frac{\theta_i}{2} + e_{i+1} e_i \sin \frac{\theta_i}{2}, \quad i = 1, \dots, n-1, \quad (11)$$

where $0 \leq \theta_1 < 2\pi$ $0 \leq \theta_i < \pi$, $i = 2, \dots, n-1$.

This follows from the description of $a \in B^n$ in spherical coordinates using the rotors (11). These rotors describe rotations in coordinate planes. For $s = \cos \left(\frac{\theta}{2} \right) + e_i e_j \sin \left(\frac{\theta}{2} \right)$, $i \neq j$ we have

$$sx\bar{s} = (\cos \theta x_i - \sin \theta x_j) e_i + (\cos \theta x_j + \sin \theta x_i) e_j + \sum_{k \neq i, j} x_k e_k,$$

which represents a rotation of angle θ in the $e_i e_j$ -plane. In general we have $s_i s_j \neq s_j s_i$, $i \neq j$. The order of the rotors is important since different choices lead to different systems of coordinates. Due to the relevance of the rotor s_{n-1} for our work we shall denote $\theta_{n-1} := \phi$.

Lemma 5 *For $a \in B^n$ as described in Lemma 4 it follows*

$$\varphi_a(x) = \varphi_{sre_n \bar{s}}(x) = s \varphi_{re_n}(\bar{s} x s) \bar{s}.$$

Combining Lemma 5 with the rotation in decomposition (8) yields the Cartan decomposition or KAK decomposition of $\text{Spin}^+(1, n)$.

For the construction of a theory of sections we need to consider some special gyro-subgroups.

Definition 3 *Let $(G, \oplus)$ be a gyrogroup and H a non-empty subset of G . H is a gyro-subgroup of $(G, \oplus)$ if it is a gyrogroup for the operation induced from G and $\text{gyr}[a, b]c \in \text{Aut}(H)$ for all $a, b, c \in H$.*

For a fixed $\omega \in S^{n-1}$, we consider the hyperplane $H_\omega = \{x \in \mathbb{R}^n : \langle \omega, x \rangle = 0\}$, the hyperdisc $D_\omega^{n-1} = H_\omega \cap B^n$ and $L_\omega = \{x \in B^n : x = t\omega, -1 < t < 1\}$. Using (3) and (1) we derive the following gyro-subgroups.

Proposition 4 *The sets D_ω^{n-1} and L_ω endowed with the operation $\oplus$ are gyro-subgroups of $(B^n, \oplus)$. Moreover, $(L_\omega, \oplus)$ is a group.*

The last property comes directly from Lemma 1. The operation $\oplus$ in L_ω corresponds to the Einstein velocity addition of parallel velocities in the special theory of relativity. The particular group $(L_{e_n}, \oplus)$ is isomorphic to the $\text{Spin}(1, 1)$ group.

4 Sections of the proper Lorentz group

We want to factorize the proper Lorentz group and to construct sections on the respective homogeneous space. First we will study the factorization of the gyrogroup $(B^n, \oplus)$ by the gyro-subgroup $(D_{e_n}^{n-1}, \oplus)$, taking into consideration only left cosets.

Theorem 2 *For each $c \in B^n$ there exist unique $a \in D_{e_n}^{n-1}$ and $b \in L_{e_n}$ such that $c = b \oplus a$.*

Proof: Let $c = (c_1, \dots, c_n) \in B^n$ be an arbitrary point. By Lemma 4 we can write $c = s_* c_* \overline{s_*}$, with $s_* = s_1 \dots s_{n-2} \in \text{Spin}(n-1)$, (this rotation leaves the x_n -axis invariant) and $c_* = (0, \dots, 0, c_{n-1}^*, c_n)$, where $c_{n-1}^* = r \sin \phi$ and $c_n = r \cos \phi$, with $r = |c| \in [0, 1[$ and $\phi = \arccos(c_n) \in [0, \pi]$.

First we prove the existence of the decomposition for c_* . If $c_{n-1}^* = 0$ then it suffices to consider $a = 0$ and $b = c_*$. If $c_{n-1}^* \neq 0$ then we consider $a = \lambda e_{n-1}$ and $b = t e_n$ such that

$$\lambda = \frac{|c_*|^2 - 1 + \sqrt{((c_n + 1)^2 + c_{n-1}^{*2})((c_n - 1)^2 + c_{n-1}^{*2})}}{2c_{n-1}^*} \quad \text{and} \quad t = \frac{c_n}{\lambda c_{n-1}^* + 1}. \quad (12)$$

We can see that $-1 < \lambda, t < 1$. Thus, $a \in D_{e_n}^{n-1}$ and $b \in L_{e_n}$. Taking into account that $a \perp b$, that is $\langle a, b \rangle = 0$, we obtain

$$b \oplus a = \left(0, \dots, 0, \frac{\lambda(1-t^2)}{1+\lambda^2 t^2}, \frac{t(1+\lambda^2)}{1+\lambda^2 t^2} \right) = c_*.$$

Now, we prove the existence result for the arbitrary point $c \in B^n$. Considering $a_* = s_* a \overline{s_*}$ then $a_* \in D_{e_n}^{n-1}$ since the rotation induced by s_* leaves the x_n -axis invariant. Thus, by (10) we have

$$b \oplus (s_* a \overline{s_*}) = s_* ((\overline{s_*} b s_*) \oplus a) \overline{s_*} = s_* (b \oplus a) \overline{s_*} = s_* c_* \overline{s_*} = c,$$

which shows that $c = b \oplus a_*$ and the existence of the decomposition is proved.

To prove the uniqueness we suppose that there exist $a, d \in D_{e_n}^{n-1}$ and $b, f \in L_{e_n}$ such that $c = b \oplus a = f \oplus d$. Then $a = (-b) \oplus (f \oplus d)$, by (6). As $b \perp d$ and $f \perp d$ we have that $a = ((-b) \oplus f) \oplus d$, by Lemma 1. Since by hypothesis $a, d \in D_{e_n}^{n-1}$, $(-b) \oplus f$ must be an element of $D_{e_n}^{n-1}$. This is true if and only if $(-b) \oplus f = 0$. This implies $b = f$ and $a = d \oplus 0 = d$. ■

If we consider the subgroup $D_{e_n}^{n-1}$ and the relation R defined by

$$\forall c, d \in B^n, \quad c R d \Leftrightarrow \exists a \in D_{e_n}^{n-1} : \quad c = d \oplus a,$$

then R is a reflexive relation but it is not symmetric nor transitive because the operation $\oplus$ is neither commutative nor associative. Thus R is not an equivalence relation. However, an equivalence relation can be built if we have a partition of the unit ball.

Proposition 5 *The family $\{S_b : b \in L_{e_n}\}$, with $S_b = \{b \oplus a : a \in D_{e_n}^{n-1}\}$ is a disjoint partition of B^n .*

Proof: We first prove that this family is indeed disjoint. Let $b = t_1 e_n$ and $c = t_2 e_n$ with $t_1 \neq t_2$, and assume that $S_b \cap S_c \neq \emptyset$. Then there exists $f \in B^n$ such that $f = b \oplus a$ and $f = c \oplus d$ for some $a, d \in D_{e_n}^{n-1}$. By (6) and (5) we have $a = (-b) \oplus (c \oplus d) = ((-b) \oplus c) \oplus (qd\bar{q})$, with $q = \frac{1+bc}{|1+bc|}$. But $q = \frac{1+bc}{|1+bc|} = \frac{1-t_1 t_2}{|1-t_1 t_2|} = 1$, hence $a = ((-b) \oplus c) \oplus d$. Then $(-b) \oplus c \in D_{e_n}^{n-1}$ because $a, d \in D_{e_n}^{n-1}$. Therefore, $(-b) \oplus c = 0$ or either $b = c$. But this contradicts our assumption. Finally, by Theorem 2 we have that $\cup_{b \in L_{e_n}} S_b = B^n$. ■

This partition induces an equivalence relation on B^n whose equivalence classes are the surfaces S_b . Two points $c, d \in B^n$ are said to be in relation if and only if there exists $b \in L_{e_n}$ such that $c, d \in S_b$, which means

$$c \sim_l d \Leftrightarrow \exists b \in L_{e_n}, \exists a, f \in D_{e_n}^{n-1} : c = b \oplus a \text{ and } d = b \oplus f. \quad (13)$$

This relation is equivalent to

$$c \sim_l d \Leftrightarrow \exists b \in L_{e_n}, \exists a, f \in D_{e_n}^{n-1} : c \oplus (q_1(-a)\bar{q}_1) = d \oplus (q_2(-f)\bar{q}_2),$$

with $q_1 = \frac{1-ab}{|1-ab|}$ and $q_2 = \frac{1-fb}{|1-fb|}$.

Thus, we have proved the isomorphism $B^n / (D_{e_n}^{n-1}, \sim_l) \cong L_{e_n}$. Given $c \in B^n$ we denote by $[c]$ its equivalence class on $B^n / (D_{e_n}^{n-1}, \sim_l)$. A natural projection can be defined by $\pi : B^n \rightarrow L_{e_n}$, such that $\pi(c) = b$, by the unique decomposition of $c \in B^n$ given in Theorem 2. More generally, we have the isomorphisms $B^n / (D_\omega^{n-1}, \sim_l) \cong L_\omega, \omega \in S^{n-1}$.

Now, we will give a characterization of the surfaces S_b .

Proposition 6 *For each $b = te_n \in L_{e_n}$ the surface S_b is a revolution surface in turn of the x_n -axis obtained by the intersection of B^n with the sphere orthogonal to S^{n-1} , with center in the point $C = (0, \dots, 0, \frac{1+t^2}{2t})$ and radius $\tau = \frac{1-t^2}{2|t|}$.*

Proof: Let $b = te_n \in L_{e_n}$, $c = \lambda e_{n-1} \in D_{e_n}^{n-1}$ and

$$P := b \oplus c = \left(0, \dots, 0, \frac{\lambda(1-t^2)}{1+\lambda^2 t^2}, \frac{t(1+\lambda^2)}{1+\lambda^2 t^2} \right).$$

Let $C_b = \{b \oplus c : c = \lambda e_{n-1}, 0 \leq \lambda < 1\}$. Each $a \in D_{e_n}^{n-1}$ can be described as $a = s_* c \bar{s}_*$, for some $c = \lambda e_{n-1}$ and $s_* = s_1 \cdots s_{n-2} \in \text{Spin}(n-1)$. Then, by (10) we have

$$b \oplus (s_* c \bar{s}_*) = s_* ((\bar{s}_* b s_*) \oplus c) \bar{s}_* = s_* (b \oplus c) \bar{s}_*.$$

Thus, S_b is a surface of revolution obtained by revolution in turn of the x_n -axis of the arc C_b . The last coordinate of the surface S_b gives the orientation of its concavity.

For all $\lambda \in [0, 1[$, we have $\|P - C\|^2 = \tau^2$, with $C = (0, \dots, 0, \frac{1+t^2}{2t})$ and $\tau = \frac{1-t^2}{2|t|}$, which means that the arc C_b lies on the sphere centered at C and radius τ . Moreover, as t tends to zero the radius of the sphere tends to infinity thus proving that the surface S_0 coincides with the hyperdisc $D_{e_n}^{n-1}$.

Each S_b is orthogonal to S^{n-1} because $\|C\|^2 = 1 + \tau^2$. We recall that two spheres, S_1 and S_2 , with centers m_1 and m_2 and radii τ_1 and τ_2 , respectively, intersect orthogonally if and only if $\langle m_1 - y, m_2 - y \rangle = 0$, for all $y \in S_1 \cap S_2$, or equivalently, if $\|m_1 - m_2\|^2 = \tau_1^2 + \tau_2^2$.

We can observe the projection of these orbits in the $x_{n-1}x_n$ -plane: ■




Figure 1: Cut of the surfaces S_b , in the $x_{n-1}x_n$ -plane.

The space $Y = B^n / (D_{e_n}^{n-1}, \sim_l)$ becomes a homogeneous space endowed with the mapping

$$h : B^n \times Y \rightarrow Y, \quad h(c, [a]) = [c \oplus a]. \quad (14)$$

We consider L_{e_n} the fundamental section σ_0 , which corresponds to the $\text{Spin}(1, 1)$ subgroup. From Proposition 6 an entire class of sections $\sigma : B^n / (D_{e_n}^{n-1}, \sim_l) \rightarrow B^n$ can be obtained from L_{e_n} by considering

$$\sigma(te_n) = te_n \oplus g(t)e_{n-1} = \left(0, \dots, 0, \frac{g(t)(1-t^2)}{1+(tg(t))^2}, \frac{t(1+g(t)^2)}{1+(tg(t))^2}\right), \quad (15)$$

where $g :]-1, 1[\rightarrow]-1, 1[$ is called the generating function of the section. Depending on the properties of the function g we can have sections that are Borel maps and also smooth sections. If $g \in C^k(]-1, 1[), k \in \mathbb{N}$, then the section generates a C^k -curve inside the unit ball. For example, for each $\lambda \in]-1, 1[$ if we consider $g(t) = \lambda, (t \in]-1, 1[)$ then we obtain the family of sections $\sigma_\lambda(t) = \left(0, \dots, 0, \frac{\lambda(1-t^2)}{1+t^2\lambda^2}, \frac{t(1+\lambda^2)}{1+t^2\lambda^2}\right)$. Another example is the family of sections $\sigma_c(\phi)$ defined for each $c \in]-1, 1[$ by $\sigma_c(\phi) = (0, \dots, 0, c \sin \phi, -\cos \phi)$, $\phi \in]0, \pi[$. The intersection of each section σ_c with the orbits S_b gives us the generating function of σ_c . For $c \in]-1, 1[\setminus \{0\}$ we obtain the generating function

$$g(t) = \begin{cases} \sqrt{\frac{t^2 - 1 + \sqrt{(1-t^2)^2 + 4c^4 t^2}}{2t^2 c^2}}, & t \in]-1, 1[\setminus \{0\} \\ c, & t = 0 \end{cases},$$

and the relation between the generating function and σ_c is given by

$$\cos \phi = \begin{cases} -\frac{t(1+g(t)^2)}{1+(tg(t))^2} = -\frac{t^2(2c^2+1)-1+\sqrt{(1-t^2)^2+4c^4 t^2}}{t(2c^2+t^2-1+\sqrt{(1-t^2)^2+4c^4 t^2})}, & t \in]-1, 1[\setminus \{0\} \\ 0, & t = 0 \end{cases}.$$

For $\lambda = 0$ or $c = 0$ we obtain the fundamental section of the homogenous space Y .

Until now we only considered the factorization of the gyrogroup $(B^n, \oplus)$. In order to incorporate rotations in our scheme we now extend the equivalence relation (13) to the group $\text{Spin}^+(1, n)$ according to the group operation (8). It is easy to see that the direct

product $\{1\} \times D_{e_n}^{n-1}$ is a gyrogroup, where 1 is the identity rotation in $\text{Spin}(n)$. Our goal is to define an equivalence relation $\sim_l^*$ on $\text{Spin}(n) \times B^n$, which is an extension of the equivalence relation $\sim_l$ on B^n , such that the homogeneous space $X = (\text{Spin}(n) \times B^n) / (\{1\} \times D_{e_n}^{n-1}, \sim_l^*)$ is isomorphic as a set to $\text{Spin}(n) \times L_{e_n}$.

For $(s_1, c), (s_2, d) \in \text{Spin}(n) \times B^n$ we define the equivalence relation $\sim_l^*$ by

$$\begin{aligned} (s_1, c) \sim_l^* (s_2, d) &\Leftrightarrow \exists s_3 \in \text{Spin}(n), \exists b \in L_{e_n}, \exists a, f \in D_{e_n}^{n-1} : \\ (s_1, c) &= \left(s_3 \frac{1 - [(-a) \oplus (b \oplus a)]a}{1 - [(-a) \oplus (b \oplus a)]a}, (-a) \oplus (b \oplus a) \right) \times (e, a) \\ &\text{and} \\ (s_2, d) &= \left(s_3 \frac{1 - [(-f) \oplus (b \oplus f)]f}{1 - [(-f) \oplus (b \oplus f)]f}, (-f) \oplus (b \oplus f) \right) \times (e, f). \end{aligned}$$

The equivalence relation (4) reduces to

$$\begin{aligned} (s_1, c) \sim_l^* (s_2, d) &\Leftrightarrow \exists s_3 \in \text{Spin}(n), \exists b \in L_{e_n}, \exists a, f \in D_{e_n}^{n-1} : \\ s_1 &= s_2 \wedge (c = b \oplus a \wedge d = b \oplus f). \end{aligned} \tag{16}$$

It is easy to see that the equivalence class associated to (s_1, c) is equal to $\{s_1\} \times [c]$, with $[c] = S_b \in B^n / (D_{e_n}^{n-1}, \sim_l)$, and the quotient space is $X = \{\{s\} \times S_b : s \in \text{Spin}(n), b \in L_{e_n}\} \cong \text{Spin}(n) \times L_{e_n}$. The group $\text{Spin}^+(1, n)$ acts on X according to (8) by the mapping

$$\begin{aligned} g : \text{Spin}^+(1, n) \times X &\rightarrow X \\ ((s_1, a), (s_2, [c])) &\mapsto \left(s_1 s_2 \frac{1 - \overline{s_2} a s_2 c}{1 - \overline{s_2} a s_2 c}, [c \oplus (\overline{s_2} a s_2)] \right). \end{aligned}$$

The space X will be the underlying homogeneous space for the construction of spherical continuous wavelet transforms.

5 Lorentz Coherent States as Wavelets on S^{n-1}

The theory of square integrability of group representations on homogeneous spaces was developed in a series of papers [4, 6] and applied for example in [14] for the case of the Weyl Heisenberg group and in [15] for Gabor analysis on the unit sphere. The theory depends only on the homogeneous space, on the choice of a section and a quasi-invariant measure on the homogeneous space, and on the representation of the group. Therefore, the general approach also works in our case since we can obtain these terms in our case. We will still use the term square-integrability modulo (H, σ) , where H denotes only a gyro-subgroup because our homogeneous space X results from the factorization of a group by a gyro-subgroup.

For the construction of a spherical continuous wavelet transform we need to define motions (rotations and translations) and dilations on S^{n-1} . Translations correspond to rotations of the homogeneous space $\text{Spin}(n)/\text{Spin}(n-1)$ and rotations can be realized as rotations around a certain axis on the sphere. Thus, both translations and rotations can be associated with the action of $\text{Spin}(n)$ on S^{n-1} . Dilations will arise from Möbius transformations of type $\varphi_a(x)$.

For $f \in L^2(S^{n-1}, dS)$, where dS is the usual Lebesgue surface measure on S^{n-1} , we define the rotation and spherical dilation operators

$$R_s f(x) = f(\overline{s}xs), \quad s \in \text{Spin}(n) \quad (17)$$

and

$$D_a f(x) = \left(\frac{1 - |a|^2}{|1 - ax|^2} \right)^{\frac{n-1}{2}} f(\varphi_{-a}(x)), \quad a \in B^n, \quad (18)$$

where $\left(\frac{1 - |a|^2}{|1 - ax|^2} \right)^{\frac{n-1}{2}}$ is the Jacobian of $\varphi_{-a}(x)$ on S^{n-1} . From these operators we construct the representation U of $\text{Spin}^+(1, n)$ in $L^2(S^{n-1}, dS)$ defined by

$$U(s, a)f(x) = R_s D_a f(x) = \left(\frac{1 - |a|^2}{|1 - a\overline{s}xs|^2} \right)^{\frac{n-1}{2}} f(\varphi_{-a}(\overline{s}xs)). \quad (19)$$

Proposition 7 $U(s, a)$ is a unitary representation of the group $\text{Spin}^+(1, n)$ in $L^2(S^{n-1}, dS)$.

Proof: We need to prove that $U(s, a)$ is a homomorphism of the group $\text{Spin}^+(1, n)$ onto the space of linear maps from the Hilbert space $L^2(S^{n-1}, dS)$ onto itself, with respect to the group operation (8).

On the one hand we have

$$\begin{aligned} U(s_1, a)U(s_2, b)f(x) &= \\ &= \left(\frac{1 - |a|^2}{|1 - a\overline{s}_1xs_1|^2} \frac{1 - |b|^2}{|1 - b\overline{s}_2\varphi_{-a}(\overline{s}_1xs_1)s_2|^2} \right)^{\frac{n-1}{2}} f(\varphi_{-b}(\overline{s}_2\varphi_{-a}(\overline{s}_1xs_1)s_2)) \\ &= \left(\frac{(1 - |a|^2)(1 - |b|^2)}{|1 - s_2b\overline{s}_2a - (a + s_2b\overline{s}_2)\overline{s}_1xs_1|^2} \right)^{\frac{n-1}{2}} f(\varphi_{-(b \oplus (\overline{s}_2as_2))}(\overline{s}_3xs_3)), \end{aligned}$$

with $s_3 = s_1s_2 \frac{1 - \overline{s}_2as_2b}{|1 - \overline{s}_2as_2b|}$.

We observe that

$$\begin{aligned} |1 - b\overline{s}_2\varphi_{-a}(\overline{s}_1xs_1)s_2|^2 &= |1 - s_2b\overline{s}_2(\overline{s}_1xs_1 + a)(1 - a\overline{s}_1xs_1)^{-1}|^2 \\ &= |1 - a\overline{s}_1xs_1|^{-2} |1 - a\overline{s}_1xs_1 - s_2b\overline{s}_2(\overline{s}_1xs_1 + a)|^2 \\ &= |1 - a\overline{s}_1xs_1|^{-2} |1 - s_2b\overline{s}_2a - (a + s_2b\overline{s}_2)\overline{s}_1xs_1|^2. \end{aligned}$$

By (9), (2) and (4) it follows

$$\begin{aligned} \varphi_{-b}(\overline{s}_2\varphi_{-a}(\overline{s}_1xs_1)s_2) &= \varphi_{-b}(\varphi_{-\overline{s}_2as_2}(\overline{s}_2\overline{s}_1xs_1s_2)) \\ &= q_1\varphi_{-(b + \overline{s}_2as_2)(1 - \overline{s}_2as_2b)^{-1}}(\overline{s}_1\overline{s}_2xs_1s_2)\overline{q_1} \\ &= \varphi_{-q_1((\overline{s}_2as_2) \oplus b)\overline{q_1}}(q_1\overline{s}_1\overline{s}_2xs_1s_2\overline{q_1}) \\ &= \varphi_{-(b \oplus (\overline{s}_2as_2))}(\overline{s}_3xs_3), \end{aligned}$$

where $q_1 = \frac{1 - b\overline{s}_2as_2}{|1 - b\overline{s}_2as_2|}$.

On the other hand we have

$$\begin{aligned} U(s_3, b \oplus (\overline{s}_2as_2))f(x) &= \left(\frac{1 - |b \oplus (\overline{s}_2as_2)|^2}{|1 - (b \oplus (\overline{s}_2as_2))\overline{s}_3xs_3|^2} \right)^{\frac{n-1}{2}} f(\varphi_{-(b \oplus (\overline{s}_2as_2))}(\overline{s}_3xs_3)) \\ &= \left(\frac{(1 - |a|^2)(1 - |b|^2)}{|1 - s_2b\overline{s}_2a - (a + s_2b\overline{s}_2)\overline{s}_1xs_1|^2} \right)^{\frac{n-1}{2}} f(\varphi_{-(b \oplus (\overline{s}_2as_2))}(\overline{s}_3xs_3)). \end{aligned}$$

We note that

$$1 - |b \oplus (\overline{s_2} a s_2)|^2 = 1 - |\varphi_{-b}(\overline{s_2} a s_2)|^2 = \frac{(1 - |a|^2)(1 - |b|^2)}{|1 - b \overline{s_2} a s_2|^2}$$

and (by (4))

$$\begin{aligned} |1 - (b \oplus (\overline{s_2} a s_2)) \overline{s_3} x s_3|^2 &= |1 - q_2(b \oplus (\overline{s_2} a s_2)) \overline{q_2} \overline{s_2} \overline{s_1} x s_1 s_2|^2 \\ &= |1 - (\overline{s_2} a s_2 \oplus b) \overline{s_2} \overline{s_1} x s_1 s_2|^2 \\ &= |1 - b \overline{s_2} a s_2|^{-2} |1 - b \overline{s_2} a s_2 - (\overline{s_2} a s_2 + b) \overline{s_2} \overline{s_1} x s_1 s_2|^2 \\ &= |1 - b \overline{s_2} a s_2|^{-2} |\overline{s_2}(1 - s_2 b \overline{s_2} a) - \overline{s_2}(a + s_2 b \overline{s_2}) \overline{s_1} x s_1|^2 \\ &= |1 - b \overline{s_2} a s_2|^{-2} |1 - s_2 b \overline{s_2} a - (a + s_2 b \overline{s_2}) \overline{s_1} x s_1|^2, \end{aligned}$$

with $q_2 = \frac{1 - \overline{s_2} a s_2 b}{|1 - \overline{s_2} a s_2 b|}$.

Thus, $U(s_1, a)(U(s_2, b)f(x)) = U(s_3, b \oplus (\overline{s_2} a s_2))f(x)$, i.e. $U(s, a)$ is a representation of $\text{Spin}^+(1, n)$ on $L^2(S^{n-1}, dS)$. As the operators R_s and D_a are unitary, so is the representation $U(s, a)$, i.e. $\|U(s, a)f\|_{L^2(S^{n-1}, dS)} = \|f\|_{L^2(S^{n-1}, dS)}$. ■

This representation can be decomposed via the Cartan decomposition of $\text{Spin}^+(1, n)$ giving rise to the well-known representations of the subgroups $K = \text{Spin}(n)$ and $A = \text{Spin}(1, 1)$ [34]. It belongs to the principal series of $SO_0(1, n)$, being irreducible on the space $L^2(S^{n-1})$.

Following the general approach of [4] we will build a system of coherent states for the Lorentz group $\text{Spin}^+(1, n)$, indexed by points of the homogeneous space X . For each section σ on $Y = B^n / (D_{e_n}^{n-1}, \sim_l)$ we will consider the parameter space $W_\sigma = \{(s, \sigma(te_n)), s \in \text{Spin}(n), t \in]-1, 1[\}$. Thus, W_σ is a section on X and we now restrict our representation U to a given section W_σ . For the fundamental section W_{σ_0} the representation $U(s, \sigma_0)$ coincides with the representation used in [9]. To see this, we only need to make the equivalence of the dilation operators. We recall that the dilation operator defined in [9] is

$$D^u \psi(\omega) = \lambda^{1/2}(u, \phi) \psi(\omega_{1/u}),$$

with $\omega = (\theta_1, \dots, \theta_{n-2}, \phi) \in S^{n-1}$, $\lambda(u, \phi) = \left(\frac{4u^2}{[(u^2-1)\cos\phi + (u^2+1)]^2} \right)^{\frac{n-1}{2}}$ and $\omega_{1/u} = ((\theta_1)_u, \dots, (\theta_{n-2})_u, \phi_u)$, with $(\theta_j)_u = \theta_j$, $j = 1, \dots, n-2$, and $\tan \frac{\phi_u}{2} = u \tan \frac{\phi}{2}$.

Proposition 8 *The dilation operators D^u and D_{te_n} are equivalent, in the sense that they produce the same action on S^{n-1} .*

Proof: The Möbius transformation φ_{te_n} corresponds to the $\text{Spin}(1, 1)$ action, which is the usual Euclidean dilation lifted on S^{n-1} by inverse stereographic projection from the South Pole [12]. It remains to show that the weights of the dilation operators D^u and D_{te_n} are equal. In fact, by the bijection between $] -1, 1[$ and $]0, \infty[$ given by $t = \frac{u-1}{u+1}$, $t \in] -1, 1[$, $u > 0$ we have

$$\left(\frac{1 - t^2}{|1 - te_n x|^2} \right)^{\frac{n-1}{2}} = \left(\frac{1 - t^2}{1 + 2 \langle te_n, x \rangle + t^2} \right)^{\frac{n-1}{2}} = \left(\frac{1 - \left(\frac{u-1}{u+1} \right)^2}{1 + 2 \frac{u-1}{u+1} x_n + \left(\frac{u-1}{u+1} \right)^2} \right)^{\frac{n-1}{2}} =$$

$$= \left(\frac{4u}{2(u^2 + 1) + 2(u^2 - 1)x_n} \right)^{\frac{n-1}{2}} = \left(\frac{4u^2}{[(u^2 - 1)\cos\phi + (u^2 + 1)]^2} \right)^{\frac{n-1}{4}}.$$

■

Next, we construct a quasi-invariant measure on X . For that, we need first to construct a quasi-invariant measure on the homogeneous space $Y = B^n / (D_{e_n}^{n-1}, \sim_l)$ and, afterwards, to extend it to the homogeneous space X .

First we observe that the measure $d\mu(te_n) = \frac{2(1-t)^{n-2}}{(1+t)^n} dt$ is a quasi-invariant measure on $Y = B^n / (D_{e_n}^{n-1}, \sim_l)$. In fact, for all $a \in B^n$ and $b \in L_{e_n}$ such that $a = s_* a_* \overline{s_*}$, with $s_* = s_1 \dots s_{n-2} \in \text{Spin}(n-1)$ (c.f. Lemma 4) we have that $-a \oplus b = (s_*(-a_*)\overline{s_*}) \oplus b = s_*(-a_* \oplus (\overline{s_*} b s_*))\overline{s_*} = s_*(-a_* \oplus b)\overline{s_*}$. Thus, the equivalence classes $[-a \oplus b]$ and $[-a_* \oplus b]$ are equal. Therefore, to prove the quasi-invariance of our measure we only need to consider the action of $a = (0, \dots, 0, a_{n-1}, a_n)$ on $b = te_n \in L_{e_n}$.

Let $d = -a \oplus b = \varphi_a(b) = \left(0, \dots, 0, \frac{-(1+t^2-2a_n t)a_{n-1}}{1-2a_n t+|a|^2 t^2}, \frac{(1-|a|^2)t-(1+t^2-2a_n t)a_n}{1-2a_n t+|a|^2 t^2}\right)$. Applying the projection formulas (12) we obtain the new equivalence class, given by

$$t^* = \tau_a(t) = \frac{-2a_n(1+t^2) + 2(1-a_{n-1}^2 + a_n^2)t}{\sqrt{C_1(t)C_2(t) + (1+|a|^2)(1+t^2) - 4ta_n}},$$

with $C_1(t) := (1-t)^2 a_{n-1}^2 + (1+t)^2 (1-a_n)^2$ and $C_2(t) := (1+t)^2 a_{n-1}^2 + (1-t)^2 (1+a_n)^2$. Differentiating with respect to t we obtain

$$\tau'_a(t) = \frac{2(1-t^2)(1+a_{n-1}^2 - a_n^2)(1-|a|^2)}{C_1(t)C_2(t) + ((1+|a|^2)(1+t^2) - 4a_n t)\sqrt{C_1(t)C_2(t)}}.$$

Therefore, the Radon-Nikodym derivative of $d\mu([-a \oplus b])$ with respect to $d\mu(b)$ is given by

$$\chi(a, b) = \frac{d\mu([-a \oplus b])}{d\mu(b)} = \frac{(1 - \tau_a(t))^{n-2} (1+t)^n}{(1 + \tau_a(t))^n (1-t)^{n-2}} \tau'_a(t).$$

For the special case of $a = a_n e_n \in L_{e_n}$ we obtain $\chi(a, b) = \left(\frac{1+a_n}{1-a_n}\right)^{n-1}$. Since for each a we have that $\tau'_a(t) > 0$, for all $t \in]-1, 1[$ we conclude that $\chi(a, b) \in \mathbb{R}^+$, for all $a \in B^n$ and $b \in L_{e_n}$, thus proving that the measure $d\mu(b)$ is quasi-invariant. For $h \in L^1(L_{e_n})$ we have

$$\int_{L_{e_n}} h([a \oplus b]) d\mu(b) = \int_{L_{e_n}} h(b) d\mu([-a \oplus b]) = \int_{L_{e_n}} h(b) \chi(a, b) d\mu(b).$$

The behavior of $\chi(a, b)$ depends on the dimension. For $n = 3$ the function $\chi(a, b)$ is bounded in the variable b , i.e. for each $a \in B^n$ there exists a constant $M(a) = \frac{(1-|a|^2)(1+a_{n-1}^2 - a_n^2)}{(1-a_n)^4} \in \mathbb{R}^+$ such that $\chi(a, b) \leq M(a)$, for all $b \in L_{e_n}$, since $\chi(a, b)$ is an increasing function of the variable b , for each $a \neq 0$. Thus, $d\mu([-a \oplus b]) \leq M(a) d\mu(b)$. For $n > 3$ this estimate is not valid since the function $\chi(a, b)$ is no longer bounded in b .

The measure $d\mu(te_n)$ is equivalent to the measure $d\mu(u) = \frac{du}{u^n}$, by means of the bijection given by $t = \frac{u-1}{u+1}$ ($u \in \mathbb{R}^+$ and $t \in]-1, 1[$). A quasi-invariant measure for X is given by $d\mu(s, te_n) = \frac{2(1-t)^{n-2}}{(1+t)^n} dt d\mu(s)$, where $d\mu(s)$ is the invariant measure on $\text{Spin}(n)$.

The measure $d\mu(s, \sigma(te_n)) = \chi(\sigma(te_n), te_n) d\mu(s, te_n)$ is the standard quasi-invariant measure for an arbitrary Borel section W_σ (see [4]). The quasi-invariant measure on a section is unique (up to equivalence), that is, if μ_1 and μ_2 are quasi-invariant measures on X then there is a Borel function $\beta : X \rightarrow \mathbb{R}^+$, such that $d\mu_1(x) = \beta(x) d\mu_2(x)$, for all $x \in X$. Thus, we will use the same measure for all Borel sections. To avoid questions of integrability we will restrict our attention only to continuous sections, that is sections $\sigma(te_n)$ such that the generating function g is of class $C([-1, 1])$.

Let $N(n, l)$ be the number of all linearly independent homogeneous harmonic polynomials of degree l in n variables and $\{Y_l^M, M = 1, \dots, N(n, l)\}_{l=0}^\infty$ be an orthonormal basis of spherical harmonics, i.e. $\langle Y_l^M, Y_{l'}^{M'} \rangle_{L^2(S^{n-1})} = \delta_{ll'} \delta_{MM'}$. Thus, a function $f \in L^2(S^{n-1})$ has the Fourier expansion

$$f(x) = \sum_{l=0}^{\infty} \sum_{M=1}^{N(n, l)} \widehat{f}(l, M) Y_l^M(x), \quad (20)$$

where $\widehat{f}(l, M) := \langle Y_l^M, f \rangle_{L^2(S^{n-1})}$ are the Fourier coefficients of f .

We will also make use of Wigner D -functions [9] which appear as the following transformation formula for spherical harmonics

$$(R(s)Y_l^M)(x) = Y_l^M(\overline{s}xs) = \sum_{M'=1}^{N(n, l)} D_{MM'}^l(s) Y_l^{M'}(x), \quad s \in \text{Spin}(n)$$

and satisfy the orthogonality relations

$$\int_{\text{Spin}(n)} \overline{D_{MN}^l(s)} D_{M'N'}^{l'}(s) d\mu(s) = \frac{1}{N(n, l)} \delta_{ll'} \delta_{MM'} \delta_{NN'}. \quad (21)$$

Using the orthogonality of spherical harmonics and Wigner-D functions we can state the following admissibility condition (c.f. Theorem 4.2, [9]).

Theorem 3 *The representation U given in (19) is square integrable modulo $(\{1\} \times D_{e_n}^{n-1}, \sim_l^*)$ and the section W_σ if there exists a nonzero admissible vector $\psi \in L^2(S^{n-1}, dS)$ satisfying*

$$\frac{1}{N(n, l)} \sum_{M=1}^{N(n, l)} \int_{-1}^1 |\widehat{\psi}_{\sigma(te_n)}(l, M)|^2 d\mu(te_n) < \infty, \quad (22)$$

uniformly in l , where $\widehat{\psi}_{\sigma(te_n)}(l, M) = \langle Y_l^M, D_{\sigma(te_n)} \psi \rangle_{L^2(S^{n-1})}$.

In order to investigate the existence of admissible functions for a given section W_σ we will project the admissibility condition on the sphere to the tangent plane. The idea is to compare every global continuous section $\sigma(te_n)$ with the fundamental section σ_0 , since by

definition every section $\sigma(te_n) = te_n \oplus f(t)e_{n-1}$ is a Möbius deformation of the fundamental section.

We consider the tangent plane of the sphere S^{n-1} at the North Pole and we denote by Φ the stereographic projection map $\Phi : S^{n-1} \rightarrow \mathbb{R}^{n-1} \cup \{\infty\}$ given by

$$\Phi(x) = \left(\frac{2x_1}{1+x_n}, \dots, \frac{2x_{n-1}}{1+x_n} \right) \quad x \in S^{n-1} \setminus \{(0, \dots, 0, -1)\},$$

and $\Phi(0, \dots, 0, -1) = \infty$. The inverse mapping $\Phi^{-1} : \mathbb{R}^{n-1} \cup \{\infty\} \rightarrow S^{n-1}$ is given by

$$\Phi^{-1}(y) = \left(\frac{4y_1}{4+|y|^2}, \dots, \frac{4y_{n-1}}{4+|y|^2}, \frac{4-|y|^2}{4+|y|^2} \right), \quad y \in \mathbb{R}^{n-1}$$

and $\Phi^{-1}(\infty) = (0, \dots, 0, -1)$.

To perform the stereographic projection of $\varphi_a(x)$ we will use the Cayley transformation in $\mathbb{R}^n$. It is a conformal mapping from the unit ball B^n into the upper half space $H_n^+ = \{x \in \mathbb{R}^n : x_n > 0\}$ defined by $y = (x + e_n)(1 + e_n x)^{-1}$. The stereographic projection Φ can be rewritten in terms of the inverse of the Cayley transformation that is

$$\Phi(x) = 2(x - e_n)(1 - e_n x)^{-1} = \left(\frac{2x_1}{1+x_n}, \dots, \frac{2x_{n-1}}{1+x_n} \right).$$

Theorem 4 *The intertwining relation $\Phi(\varphi_a(x)) = \tilde{\varphi}(\Phi(x))$ holds, where $\tilde{\varphi}(x)$ is the Möbius transformation in $\mathbb{R}^{n-1}$ defined by the Vahlen matrix*

$$\frac{1}{\sqrt{1-|a|^2}} \begin{pmatrix} 1+a_n & 2(-a+a_n e_n) \\ \frac{a-a_n e_n}{2} & 1-a_n \end{pmatrix}. \quad (23)$$

Proof: Solving the system of equations in order to $u, v, w, z \in \Gamma(n)$ we obtain:

$$\begin{pmatrix} 2 & -2e_n \\ -e_n & 1 \end{pmatrix} \begin{pmatrix} 1 & -a \\ a & 1 \end{pmatrix} = \begin{pmatrix} u & v \\ w & z \end{pmatrix} \begin{pmatrix} 2 & -2e_n \\ -e_n & 1 \end{pmatrix} \\ \Leftrightarrow \begin{cases} 2u - ve_n = 2 - 2e_n a \\ -2ue_n + v = -2a - 2e_n \\ 2w - ze_n = -e_n + a \\ -2we_n + z = 1 + e_n a \end{cases} \Leftrightarrow \begin{cases} u = \frac{2-2e_n a + ve_n}{2} \\ v = -2a - 2e_n + 2ue_n \\ w = \frac{-e_n + a + ze_n}{2} \\ z = 1 + e_n a + 2we_n \end{cases} \Leftrightarrow \begin{cases} u = 1 + a_n \\ v = -2a + 2a_n e_n \\ w = \frac{a-a_n e_n}{2} \\ z = 1 - a_n \end{cases}.$$

In the resolution of the system we used the identities $e_n a e_n = -2a_n e_n + a$ and $e_n a + a e_n = -2a_n$. After normalization we obtain the matrix (23), which is in the Clifford group $\Gamma(1, n+1)$, by Theorem 1, and thus, it represents a Möbius transformation in $\mathbb{R}^{n-1}$. ■

Now, we will consider the unitary stereographic projection operator Θ from $L^2(S^{n-1}, dS)$ into $L^2(\mathbb{R}^{n-1}, r^{n-2} dr dS^{n-2})$ and we will find the stereographic mapping in $\mathbb{R}^{n-1}$ that intertwines with the dilation operator D_a . To distinguish functions from different spaces we will use the convention $f \in L^2(S^{n-1})$ and $F \in L^2(\mathbb{R}^{n-1})$.

Lemma 6 *The map $\Theta : L^2(S^{n-1}, dS) \rightarrow L^2(\mathbb{R}^{n-1}, r^{n-2} dr dS^{n-2})$ defined by*

$$f(\theta_1, \dots, \theta_{n-2}, \phi) \mapsto F(\theta_1, \dots, \theta_{n-2}, r) = \left(\frac{4}{4+r^2} \right)^{\frac{n-1}{2}} f(\theta_1, \dots, \theta_{n-2}, 2 \arctan(r/2))$$

is a unitary map. In cartesian coordinates it reads as

$$f(x) \mapsto F(y) = \left(\frac{4}{4+|y|^2} \right)^{\frac{n-1}{2}} f(\Phi_1^{-1}(y)). \quad (24)$$

Proof: By the change of variables $\phi = 2 \arctan(r/2)$, that is, $r = 2 \tan(\phi/2) = 2\sqrt{\frac{1-\cos\phi}{1+\cos\phi}}$ we have

$$\begin{aligned} \|\Theta f\|^2 &= \int_{S^{n-2}} \int_0^\infty \left| \left(\frac{4}{4+r^2} \right)^{\frac{n-1}{2}} f(\theta_1, \dots, \theta_{n-2}, 2 \arctan(r/2)) \right|^2 r^{n-2} dr dS^{n-2} \\ &= \int_{S^{n-2}} \int_0^\pi |f(\theta_1, \dots, \theta_{n-2}, \phi)|^2 \sin^{n-2} \phi d\phi dS^{n-2} \\ &= \|f\|_{L^2(S^{n-1}, dS)}^2. \end{aligned}$$

■

Theorem 5 *Let $y \in \mathbb{R}^{n-1}$, and $\tilde{\varphi}(y) := \frac{c_1 y + c_2}{c_3 y + c_4}$ be the Möbius transformation obtained from the matrix (23), with $c_1 = \frac{1+a_n}{\sqrt{1-|a|^2}}$, $c_2 = \frac{2(-a+a_n e_n)}{\sqrt{1-|a|^2}}$, $c_3 = \frac{a-a_n e_n}{2\sqrt{1-|a|^2}}$ and $c_4 = \frac{1-a_n}{\sqrt{1-|a|^2}}$. Then we have the intertwining relation*

$$\Theta D_a \psi = M \Theta \psi, \quad (25)$$

where $MF(y) = \left(\frac{4(1-|a|^2)}{|-(a-a_n e_n)y + 2(1+a_n)|^2} \right)^{\frac{n-1}{2}} F(\tilde{\varphi}^{-1}(y))$ is the unitary operator associated with $\tilde{\varphi}^{-1}(y) = \frac{c_4 y - c_2}{-c_3 y + c_1}$.

Proof: By definition we have

$$\Theta D_a \psi(y) = \left(\frac{4}{4+|y|^2} \right)^{\frac{n-1}{2}} \left(\frac{1-|a|^2}{|1-a\Phi^{-1}(y)|^2} \right)^{\frac{n-1}{2}} \psi(\varphi_{-a}(\Phi_1^{-1}(y))) \quad (26)$$

and

$$M \Theta \psi(y) = \left(\frac{4(1-|a|^2)}{|-(a-a_n e_n)y + 2(1+a_n)|^2} \right)^{\frac{n-1}{2}} \left(\frac{4}{4+|\tilde{\varphi}^{-1}(y)|^2} \right)^{\frac{n-1}{2}} \psi(\Phi_1^{-1}(\tilde{\varphi}^{-1}(y))). \quad (27)$$

First we observe that

$$\varphi_{-a}(\Phi^{-1}(y)) = \Phi^{-1}(\tilde{\varphi}^{-1}(y)), \quad \forall y \in \overline{\mathbb{R}^{n-1}}. \quad (28)$$

If $\varphi_{-a}(\Phi^{-1}(y)) = x \in S^{n-1}$ then $y = \Phi(\varphi_a(x))$. Moreover, if $\Phi^{-1}(\tilde{\varphi}^{-1}(y)) = x$, then $y = \tilde{\varphi}(\Phi(x))$. Since the relation $\Phi(\varphi_a(x)) = \tilde{\varphi}(\Phi(x))$ holds (Theorem 4) we conclude that

the relation (28) is true. It remains to prove that the weights in (26) and (27) are equal. On the one hand,

$$\begin{aligned} \frac{4}{4+|y|^2} \frac{1-|a|^2}{|1-a\Phi^{-1}(y)|^2} &= \frac{4}{4+|y|^2} \frac{1-|a|^2}{1+2\langle a, \Phi^{-1}(y) \rangle + |a|^2} \\ &= \frac{4(1-|a|^2)}{(1+|a|^2)(4+|y|^2) + 2a_n(4-|y|^2) + 8(\langle a, y \rangle - a_n y_n)}. \end{aligned}$$

On the other hand, since

$$|\tilde{\varphi}^{-1}(y)|^2 = \frac{|c_4 y - c_2|^2}{|-c_3 y + c_1|^2} = 4 \frac{(1-a_n)^2 |y|^2 + 4(1-a_n)(\langle a, y \rangle - a_n y_n) + 4(|a|^2 - a_n^2)}{(|a|^2 - a_n^2)|y|^2 + 4(1+a_n)(\langle a, y \rangle - a_n y_n) + 4(1+a_n)^2}$$

and

$$|-(a - a_n e_n)y + 2(1 + a_n)|^2 = (|a|^2 - a_n^2)|y|^2 + 4(1 + a_n)(\langle a, y \rangle - a_n y_n) + 4(1 + a_n)^2$$

we obtain, after some computations

$$\begin{aligned} \frac{4(1-|a|^2)}{|-(a - a_n e_n)y + 2(1 + a_n)|^2} \frac{4}{4+|\tilde{\varphi}^{-1}(y)|^2} &= \\ \frac{4(1-|a|^2)}{(1+|a|^2)(4+|y|^2) + 2a_n(4-|y|^2) + 8(\langle a, y \rangle - a_n y_n)}. \end{aligned}$$

Thus, the result follows. ■

The group $SL_2(\Gamma(n) \cup \{0\})$ admits an Iwasawa decomposition similar to the decomposition of the group $SL_2(\mathbb{C})$. The Iwasawa decomposition of a generic element of $SL_2(\Gamma(n) \cup \{0\})$ is

$$\begin{pmatrix} u & v \\ w & z \end{pmatrix} = \begin{pmatrix} \alpha & \beta \\ -\bar{\beta} & \bar{\alpha} \end{pmatrix} \begin{pmatrix} \delta^{-1/2} & 0 \\ 0 & \delta^{1/2} \end{pmatrix} \begin{pmatrix} 1 & \xi \\ 0 & 1 \end{pmatrix} \quad (29)$$

where $\alpha, \beta, \xi \in \Gamma(n) \cup \{0\}$ and $\delta \in \mathbb{R}^+$, $uz^* - vw^* = 1$ and

$$\delta = (|u|^2 + |w|^2)^{-1}, \quad \alpha = u\delta^{1/2}, \quad \beta = -\bar{w}\delta^{1/2},$$

$$\xi = u^{-1}(v + \bar{w}\delta), \text{ if } u \neq 0 \quad \text{or} \quad \xi = w^{-1}(z - \bar{u}\delta), \text{ if } w \neq 0.$$

The Iwasawa decomposition of the matrix 23 yields the parameters

$$\begin{aligned} \alpha &= \frac{2(1+a_n)}{\sqrt{4(1+a_n)^2 + |a|^2 - a_n^2}}, & \beta &= \frac{a - a_n e_n}{\sqrt{4(1+a_n)^2 + |a|^2 - a_n^2}}, \\ \delta &= \frac{4(1-|a|^2)}{4(1+a_n)^2 + |a|^2 - a_n^2}, & \xi &= \frac{2(-a + a_n e_n)(5 + 3a_n)}{4(1+a_n)^2 + |a|^2 - a_n^2}. \end{aligned} \quad (30)$$

Departing from the Iwasawa decomposition with parameters (30) we define the following unitary operators on $L^2(\mathbb{R}^{n-1})$:

$$R^{\alpha, \beta} F(y) = \left(\frac{1}{|-\beta \bar{y} + \alpha|^2} \right)^{\frac{n-1}{2}} F\left(\frac{\alpha y + \beta}{-\beta \bar{y} + \alpha} \right);$$

$$D^\delta F(y) = \delta^{-\frac{n-1}{2}} F\left(\frac{y}{\delta}\right); \quad T^\xi F(y) = F(y + \xi).$$

As these mappings are isometries whose ranges include the whole space $L^2(\mathbb{R}^{n-1})$ the adjoint and inverse operators are identical.

Lemma 7 *The adjoint operators to $R^{\alpha,\beta}$, D^δ and T^ξ are given by*

$$(R^{\alpha,\beta})^* = R^{\bar{\alpha},-\beta}, \quad (D^\delta)^* = D^{\frac{1}{\delta}}, \quad \text{and} \quad (T^\xi)^* = T^{-\xi}. \quad (31)$$

Proposition 9 *The operator M admits the following factorization*

$$MF = R^{\alpha,-\beta} D^{\frac{1}{\delta}} T^{-\xi} F. \quad (32)$$

Proof: We have

$$MF(y) = \left(\frac{4(1-|a|^2)}{|-(a-a_n e_n)y + 2(1+a_n)|^2} \right)^{\frac{n-1}{2}} F(\tilde{\varphi}^{-1}(y))$$

and

$$R^{\alpha,-\beta} D^{\frac{1}{\delta}} T^{-\xi} F(y) = \left(\frac{\delta}{|\alpha - \beta y|^2} \right)^{\frac{n-1}{2}} F\left(\delta \frac{\alpha y - \beta}{\beta y + \alpha} - \xi\right).$$

First we prove the equality between arguments. On the one hand,

$$\tilde{\varphi}^{-1}(y) = \frac{c_4 y - c_2}{-c_3 y + c_1} = \frac{2(1-a_n)y - 4(-a + a_n e_n)}{(-a + a_n e_n)y + 2(1+a_n)}.$$

On the other hand,

$$\delta \frac{\alpha y - \beta}{\beta y + \alpha} - \xi = \frac{(\delta\alpha - \bar{\beta}\xi)y - (\delta\beta + \alpha\xi)}{\bar{\beta}y + \alpha} = \frac{2(1-a_n)y - 4(-a + a_n e_n)}{(-a + a_n e_n)y + 2(1+a_n)}.$$

Finally, since $\left(\frac{\delta}{|\alpha - \beta y|^2}\right)^{\frac{n-1}{2}} = \left(\frac{4(1-|a|^2)}{|-(a-a_n e_n)y + 2(1+a_n)|^2}\right)^{\frac{n-1}{2}}$ we have the desired factorization. ■

Corollary 2 *The intertwining relation (25) can be written as*

$$\Theta D_a \psi = R^{\alpha,-\beta} D^{\frac{1}{\delta}} T^{-\xi} \Theta \psi. \quad (33)$$

Corollary 3 *For $a = te_n \in L_{e_n}$ (restriction to the $Spin(1,1)$ case) we obtain the well-known intertwining relation*

$$\Theta D_{te_n} \psi = D^{\frac{1+t}{1-t}} \Theta \psi. \quad (34)$$

Relation (33) is the generalization of relation (34). The first case correspond to the isotropic case of Antoine and vandergheynst and the second case correspond to the anisotropic case, for $a \in B^n \setminus L_{e_n}$.

For an arbitrary global left section $\sigma(te_n) = te_n \oplus g(t)e_{n-1}$ (15), the parameters (30) become

$$\begin{aligned}\alpha_t &= \frac{2(1+tg(t)^2)}{\sqrt{(1+t^2+6t)g(t)^2+4(1+t^2g(t)^4)}}, \\ \beta_t &= \frac{(1-t)g(t)}{\sqrt{(1+t^2+6t)g(t)^2+4(1+t^2g(t)^4)}} e_{n-1}, \\ \delta_t &= \frac{4(1-t)(1-g(t)^2)(1+t^2g(t)^2)}{(t+1)((1+t^2+6t)g(t)^2+4(1+t^2g(t)^4))}, \\ \xi_t &= \frac{2(t-1)g(t)((5t^2+3t)g(t)^2+3t+5)}{(t+1)((1+t^2+6t)g(t)^2+4(1+t^2g(t)^4))} e_{n-1}.\end{aligned}\tag{35}$$

When $g(t) \equiv 0$, (restriction to the fundamental section - Spin(1,1) case) we obtain $\alpha = 1$, $\beta = 0$, $\xi = 0$ and $\delta = \frac{1-t}{1+t}$, which reflects again the fact that we obtain a pure dilation.

Lemma 8 *The parameter δ_t can be written as a deformation of the pure dilation parameter, i.e. $\delta_t = \frac{1-t}{1+t}\delta_t^*$, with $\delta_t^* = \frac{4(1-g(t)^2)(1+t^2g(t)^2)}{(1+t^2+6t)g(t)^2+4(1+t^2g(t)^4)}$, and where δ_t^* satisfies the estimate $0 < \delta_t^* < \frac{2(3-2\sqrt{3})}{3(-2+\sqrt{3})}$, for all $t \in]-1, 1[$.*

Proof: As $g(t) \in]-1, 1[$, for every $t \in]-1, 1[$, the study of the behavior of the parameter δ_t^* is equivalent to the study of the behavior of the function of two variables $h(t, \lambda) = \frac{4(1-\lambda^2)(1+t^2\lambda^2)}{(1+t^2+6t)\lambda^2+4(1+t^2\lambda^4)}$, with $t, \lambda \in]-1, 1[$. Since for each $\lambda \in]-1, 1[$

$$\frac{1-\lambda^2}{1+\lambda^2} < h(t, \lambda) < \frac{1-\lambda^4}{1+\lambda^4-\lambda^2}, \quad \forall t \in]-1, 1[$$

we conclude that

$$0 < h(t, \lambda) < \frac{2(3-2\sqrt{3})}{3(-2+\sqrt{3})}, \quad \forall t, \lambda \in]-1, 1[.$$

Therefore,

$$0 < \delta_t^* < \frac{2(3-2\sqrt{3})}{3(-2+\sqrt{3})}, \quad \forall t \in]-1, 1[.$$

■

Now, we will prove the main theorem of this paper concerning the square-integrability of the representation U over an arbitrary continuous section W_σ .

Theorem 6 *Let $\psi \in L^2(S^{n-1})$ such that the family $\{R_s D_{te_n} \psi, s \in \text{Spin}(n), t \in]-1, 1[\}$ is a continuous frame, that is, there exist constants $0 < A \leq B < \infty$ such that*

$$A\|f\|^2 \leq \int_{\text{Spin}(n)} \int_{-1}^1 |\langle f, R_s D_{te_n} \psi \rangle|^2 d\mu(te_n) d\mu(s) \leq B\|f\|^2, \quad \forall f \in L^2(S^{n-1}). \tag{36}$$

Then ψ is an admissible function for any continuous section W_σ and the system $\{R_s D_{\sigma(te_n)} \psi, s \in \text{Spin}(n), t \in]-1, 1[\}$ is also a continuous frame.

Proof: For every $a \in B^n$ and $f \in L^2(S^{n-1})$ arbitrary we have

$$\begin{aligned}
& \int_{\text{Spin}(n)} \int_{-1}^1 \left| \langle f, R_s D_a \psi \rangle_{L^2(S^2)} \right|^2 d\mu(te_n) d\mu(s) \\
&= \int_{\text{Spin}(n)} \int_{-1}^1 \left| \langle R_{\bar{s}} f, D_a \psi \rangle_{L^2(S^2)} \right|^2 d\mu(te_n) d\mu(s) \\
&= \int_{\text{Spin}(n)} \int_{-1}^1 \left| \langle \Theta R_{\bar{s}} f, \Theta D_a \psi \rangle_{L^2(\mathbb{R}^2)} \right|^2 d\mu(te_n) d\mu(s) \quad (\text{by Lemma 6}) \\
&= \int_{\text{Spin}(n)} \int_{-1}^1 \left| \left\langle \Theta R_{\bar{s}} f, R^{\alpha, -\beta} D^{\frac{1}{\delta}} T^{-\xi} \Theta \psi \right\rangle_{L^2(\mathbb{R}^{n-1})} \right|^2 d\mu(te_n) d\mu(s) \quad (\text{by (33)}) \\
&= \int_{\text{Spin}(n)} \int_{-1}^1 \left| \left\langle \Theta R_{\bar{s}} f, R^{\alpha, -\beta} T^{-\frac{\xi}{\delta}} D^{\frac{1}{\delta}} \Theta \psi \right\rangle_{L^2(\mathbb{R}^{n-1})} \right|^2 d\mu(te_n) d\mu(s) \\
&= \int_{\text{Spin}(n)} \int_{-1}^1 \left| \left\langle T^{\frac{\xi}{\delta}} R^{\alpha, \beta} \Theta R_{\bar{s}} f, D^{\frac{1}{\delta}} \Theta \psi \right\rangle_{L^2(\mathbb{R}^{n-1})} \right|^2 d\mu(te_n) d\mu(s) \quad (\text{by (7)}). \tag{37}
\end{aligned}$$

Now we consider $a = \sigma(te_n)$ and the parameters (35). By Lemma 8, the integral (37) becomes

$$\int_{\text{Spin}(n)} \int_{-1}^1 \left| \left\langle D^{\delta_t^*} T^{\frac{\xi_t}{\delta_t}} R^{\alpha_t, \beta_t} \Theta R_{\bar{s}} f, D^{\frac{1+t}{1-t}} \Theta \psi \right\rangle_{L^2(\mathbb{R}^{n-1})} \right|^2 d\mu(te_n) d\mu(s). \tag{38}$$

For each $s \in \text{Spin}(n)$ and $t \in]-1, 1[$ we have

$$\left| \left\langle D^{\delta_t^*} T^{\frac{\xi_t}{\delta_t}} R^{\alpha_t, \beta_t} \Theta R_{\bar{s}} f, D^{\frac{1+t}{1-t}} \Theta \psi \right\rangle \right|^2 \leq \underbrace{\sup_{t' \in]-1, 1[} \left| \left\langle D^{\delta_{t'}^*} T^{\frac{\xi_{t'}}{\delta_{t'}}} R^{\alpha_{t'}, \beta_{t'}} \Theta R_{\bar{s}} f, D^{\frac{1+t}{1-t}} \Theta \psi \right\rangle \right|^2}_I$$

and

$$\underbrace{\inf_{t' \in]-1, 1[} \left| \left\langle D^{\delta_{t'}^*} T^{\frac{\xi_{t'}}{\delta_{t'}}} R^{\alpha_{t'}, \beta_{t'}} \Theta R_{\bar{s}} f, D^{\frac{1+t}{1-t}} \Theta \psi \right\rangle \right|^2}_{II} \leq \left| \left\langle D^{\delta_t^*} T^{\frac{\xi_t}{\delta_t}} R^{\alpha_t, \beta_t} \Theta R_{\bar{s}} f, D^{\frac{1+t}{1-t}} \Theta \psi \right\rangle \right|^2.$$

Let $t_1, t_2 \in [-1, 1]$ such that

$$I := \left| \left\langle D^{\delta_{t_2}^*} T^{\frac{\xi_{t_2}}{\delta_{t_2}}} R^{\alpha_{t_2}, \beta_{t_2}} \Theta R_{\bar{s}} f, D^{\frac{1+t}{1-t}} \Theta \psi \right\rangle \right|^2$$

and

$$II := \left| \left\langle D^{\delta_{t_1}^*} T^{\frac{\xi_{t_1}}{\delta_{t_1}}} R^{\alpha_{t_1}, \beta_{t_1}} \Theta R_{\bar{s}} f, D^{\frac{1+t}{1-t}} \Theta \psi \right\rangle \right|^2.$$

Thus, we obtain

$$\begin{aligned}
& \int_{\text{Spin}(n)} \int_{-1}^1 \left| \left\langle D_{t_1}^{\delta_{t_1}^*} T_{\delta_{t_1}}^{\frac{\xi_{t_1}}{\delta_{t_1}}} R^{\alpha_{t_1}, \beta_{t_1}} \Theta R_{\bar{s}} f, D^{\frac{1+t}{1-t}} \Theta \psi \right\rangle_{L^2(\mathbb{R}^{n-1})} \right|^2 d\mu(te_n) d\mu(s) \leq \\
& \int_{\text{Spin}(n)} \int_{-1}^1 \left| \left\langle D_{t_2}^{\delta_{t_2}^*} T_{\delta_{t_2}}^{\frac{\xi_{t_2}}{\delta_{t_2}}} R^{\alpha_{t_2}, \beta_{t_2}} \Theta R_{\bar{s}} f, D^{\frac{1+t}{1-t}} \Theta \psi \right\rangle_{L^2(\mathbb{R}^{n-1})} \right|^2 d\mu(te_n) d\mu(s) \leq \\
& \int_{\text{Spin}(n)} \int_{-1}^1 \left| \left\langle D_{t_1}^{\delta_{t_1}^*} T_{\delta_{t_1}}^{\frac{\xi_{t_1}}{\delta_{t_1}}} R^{\alpha_{t_1}, \beta_{t_1}} \Theta R_{\bar{s}} f, D^{\frac{1+t}{1-t}} \Theta \psi \right\rangle_{L^2(\mathbb{R}^{n-1})} \right|^2 d\mu(te_n) d\mu(s).
\end{aligned}$$

Since for $t \in [-1 + \epsilon, 1 - \epsilon]$, $\epsilon > 0$, the operators $D_t^{\delta_t^*}$, $T_{\delta_t}^{\frac{\xi_t}{\delta_t}}$ and R^{α_t, β_t} are unitary and bijective mappings we only need to study the case of $t \rightarrow \pm 1$. In this case, the parameter $\frac{\xi_t}{\delta_t}$ associated to the operator $T_{\delta_t}^{\frac{\xi_t}{\delta_t}}$ can become infinity. But, it is easy to see that the composition of the operators $D_t^{\delta_t^*}$ and $T_{\delta_t}^{\frac{\xi_t}{\delta_t}}$ is well behaved. Thus, for each $f \in L^2(S^{n-1})$ we can find $f_1, f_2 \in L^2(S^{n-1})$ such that $D_{t_1}^{\delta_{t_1}^*} T_{\delta_{t_1}}^{\frac{\xi_{t_1}}{\delta_{t_1}}} R^{\alpha_{t_1}, \beta_{t_1}} \Theta R_{\bar{s}} f = \Theta R_{\bar{s}} f_1$ and $D_{t_2}^{\delta_{t_2}^*} T_{\delta_{t_2}}^{\frac{\xi_{t_2}}{\delta_{t_2}}} R^{\alpha_{t_2}, \beta_{t_2}} \Theta R_{\bar{s}} f = \Theta R_{\bar{s}} f_2$ with $\|f_1\| = \|f_2\| = \|f\|$.

Therefore, we have

$$\begin{aligned}
& \int_{\text{Spin}(n)} \int_{-1}^1 \left| \left\langle \Theta R_{\bar{s}} f_1, D^{\frac{1+t}{1-t}} \Theta \psi \right\rangle_{L^2(\mathbb{R}^{n-1})} \right|^2 d\mu(te_n) d\mu(s) \leq \\
& \int_{\text{Spin}(n)} \int_{-1}^1 \left| \left\langle D_{t_1}^{\delta_{t_1}^*} T_{\delta_{t_1}}^{\frac{\xi_{t_1}}{\delta_{t_1}}} R^{\alpha_{t_1}, \beta_{t_1}} \Theta R_{\bar{s}} f, D^{\frac{1+t}{1-t}} \Theta \psi \right\rangle_{L^2(\mathbb{R}^{n-1})} \right|^2 d\mu(te_n) d\mu(s) \leq \\
& \int_{\text{Spin}(n)} \int_{-1}^1 \left| \left\langle \Theta R_{\bar{s}} f_2, D^{\frac{1+t}{1-t}} \Theta \psi \right\rangle_{L^2(\mathbb{R}^{n-1})} \right|^2 d\mu(te_n) d\mu(s). \tag{39}
\end{aligned}$$

By (33), (37), and Lemma 6, condition (39) becomes

$$\begin{aligned}
& \int_{\text{Spin}(n)} \int_{-1}^1 \left| \langle f_1, R_s D_{te_n} \psi \rangle_{L^2(S^{n-1})} \right|^2 d\mu(te_n) d\mu(s) \leq \\
& \int_{\text{Spin}(n)} \int_{-1}^1 \left| \langle f, R_s D_{\sigma(te_n)} \psi \rangle_{L^2(S^{n-1})} \right|^2 d\mu(te_n) d\mu(s) \leq \\
& \int_{\text{Spin}(n)} \int_{-1}^1 \left| \langle f_2, D_{te_n} \Theta \psi \rangle_{L^2(S^{n-1})} \right|^2 d\mu(te_n) d\mu(s). \tag{40}
\end{aligned}$$

As, by hypothesis, ψ satisfies condition (36) there exist constants $0 < A \leq B < \infty$ such that

$$A\|f_1\|^2 \leq \int_{\text{Spin}(n)} \int_{-1}^1 |\langle f, R_s D_{\sigma(te_n)} \psi \rangle|^2 d\mu(te_n) d\mu(s) \leq B\|f_2\|^2, \quad \forall f \in L^2(S^{n-1}),$$

and so, since $\|f_1\| = \|f_2\| = \|f\|$, the result holds.

As a consequence of this theorem we conclude that there exists $\psi \in L^2(S^{n-1})$ such that the condition (22) is satisfied. ■

For an arbitrary section W_σ and an admissible function $\psi \in L^2(S^{n-1})$ we define the **generalized spherical CWT**

$$W_\psi[f](s, \sigma(te_n)) = \langle R_s D_{\sigma(te_n)} \psi, f \rangle = \int_{S^{n-1}} \overline{R_s D_{\sigma(te_n)} \psi(x)} f(x) dS. \quad (41)$$

For each section σ and each function $\psi \neq 0$ we define an operator A_σ^ψ by

$$\langle A_\sigma^\psi f, f \rangle = \int_{\text{Spin}(n)} \int_{-1}^1 |\langle f, R_s D_{\sigma(te_n)} \psi \rangle|^2 d\mu(te_n) d\mu(s).$$

It is a Fourier multiplier given by $\widehat{A_\sigma^\psi h}(l, M) = C_\sigma^\psi(l) \widehat{h}(l, M)$, with

$$C_\sigma^\psi(l) := \frac{1}{N(n, l)} \sum_{M=1}^{N(n, l)} \int_{-1}^1 |\widehat{\psi}_{\sigma(te_n)}(l, M)|^2 d\mu(te_n).$$

By Theorem 6, for an admissible wavelet the operator A_σ is bounded with bounded inverse, since there exist constants $0 < C_1 \leq C_2 < \infty$ such that $C_1 \leq C_\sigma^\psi(l) \leq C_2$.

The wavelet transform (41) is a mapping from $L^2(S^{n-1}, dS)$ to $L^2(\text{Spin}(n) \times \sigma(te_n), d\mu(s) d\mu(te_n))$. Moreover, there exists a reconstruction formula and also a Plancherel Theorem.

Proposition 10 *Let $f \in L^2(S^{n-1})$. If ψ is an admissible wavelet then*

$$f(x) = \int_{-1}^1 \int_{\text{Spin}(n)} W_\psi[f](s, \sigma(te_n)) [R_s (A_\sigma^\psi)^{-1} D_{\sigma(te_n)} \psi](x) d\mu(s) d\mu(te_n). \quad (42)$$

Proof: Let $\psi_{\sigma(te_n)} := D_{\sigma(te_n)} \psi$. Then we have

$$[R_s (A_\sigma^\psi)^{-1} \psi_{\sigma(te_n)}](x) = \sum_{l \in \mathbb{N}} \sum_M \sum_N \frac{1}{C_\sigma^\psi(l)} D_{MN}^l(s) \widehat{\psi}_{\sigma(te_n)}(l, M) Y_l^N(x). \quad (43)$$

The wavelet coefficients $W_\psi[f](s, \sigma(te_n))$ defined in (41) may be written as

$$W_\psi[f](s, \sigma(te_n)) = \sum_{l \in \mathbb{N}} \sum_M \sum_N \overline{D_{MN}^l(s) \widehat{\psi}_{\sigma(te_n)}(l, M)} \widehat{f}(l, N). \quad (44)$$

Inserting expressions (43) and (44) in (42) and using the orthogonality relation for Wigner

D -functions, we obtain

$$\begin{aligned}
& \int_{-1}^1 \int_{\text{Spin}(n)} W_\psi[f](s, \sigma(te_n)) [R_s(A_\sigma^\psi)^{-1} D_{\sigma(te_n)} \psi](x) d\mu(s) d\mu(te_n) = \\
& = \int_{-1}^1 \int_{\text{Spin}(n)} \sum_{l \in \mathbb{N}} \sum_M \sum_N \overline{D_{MN}^l(s)} \widehat{\psi}_{\sigma(te_n)}(l, M) \widehat{f}(l, N) \\
& \quad \sum_{l' \in \mathbb{N}} \sum_{M'} \sum_{N'} \frac{1}{C_\sigma^\psi(l')} D_{M'N'}^{l'}(s) \widehat{\psi}_{\sigma(te_n)}(l', M') Y_{l'}^{N'}(x) d\mu(s) d\mu(te_n) \\
& = \sum_{l, l'} \sum_{M, M'} \sum_{N, N'} \int_{-1}^1 \overline{\widehat{\psi}_{\sigma(te_n)}(l, M)} \widehat{\psi}_{\sigma(te_n)}(l', M') d\mu(te_n) \widehat{f}(l, N) Y_{l'}^{N'}(x) \\
& \quad \frac{1}{C_\sigma^\psi(l')} \int_{\text{Spin}(n)} \overline{D_{MN}^l(s)} D_{M'N'}^{l'}(s) d\mu(s) \\
& = \sum_{l \in \mathbb{N}} \sum_N \widehat{f}(l, N) Y_l^N(x) \frac{1}{C_\sigma^\psi(l)} \left[\frac{1}{N(n, l)} \sum_M \int_{-1}^1 |\widehat{\psi}_{\sigma(te_n)}(l, M)|^2 d\mu(te_n) \right] \quad (\text{by (21)}) \\
& = \sum_{l \in \mathbb{N}} \sum_N \widehat{f}(l, N) Y_l^N(x) \\
& = f(x).
\end{aligned}$$

■

Corollary 4 *For an admissible wavelet ψ the following Plancherel relation is satisfied*

$$||f||^2 = \int_{-1}^1 \int_{\text{Spin}(n)} \overline{\widetilde{W}_\psi[f](s, \sigma(te_n))} W_\psi[f](s, \sigma(te_n)) d\mu(s) d\mu(te_n) \quad (45)$$

with

$$\widetilde{W}_\psi[f](s, \sigma(te_n)) = \left\langle \widetilde{\psi}_{(s, \sigma(te_n))}, f \right\rangle = \left\langle R_s(A_\sigma^\psi)^{-1} D_{\sigma(te_n)} \psi, f \right\rangle. \quad (46)$$

As a consequence of Theorem 6, every admissible function for the fundamental section W_{σ_0} is also an admissible function for any global continuous section W_σ . We propose the following definition for the new wavelets arising from the action of the conformal group.

Definition 4 *For a given global continuous section W_σ and an admissible $\psi \in L^2(S^{n-1})$ the wavelets (or system of coherent states) obtained as*

$$\psi_{(s, \sigma(te_n))}(x) := U(s, \sigma(te_n))\psi(x) = R_s D_{\sigma(te_n)} \psi, \quad s \in \text{Spin}(n), t \in]-1, 1[$$

are called **spherical conformlets**.

6 Anisotropy and covariance of the generalized SCWT

The generalized spherical CWT depends on the chosen section, where the pure dilation operator is replaced by an anisotropic dilation operator varying continuously from scale to

scale. One essential question is to understand what kind of dilations are obtained when the Möbius transformation $\varphi_{te_n}(x)$ is replaced by a Möbius transformation $\varphi_{\sigma(te_n)(x)}$. The study made in [12] of the influence of the parameter $a \in B^n$ on a given spherical cap centered at the North Pole allow us to understand the local character of anisotropic dilations. By the Unique Decomposition Theorem (Theorem 2) we obtain the following factorization of the dilation operator D_c , for each $c \in B^n$.

Proposition 11 *Let $c \in B^n$ such that $c = b \oplus a$, with $a \in D_{e_n}^{n-1}$ and $b \in L_{e_n}$. Then*

$$D_c f(x) = D_b D_a R_q f(x), \quad \text{with} \quad q = \frac{1 - ab}{|1 - ab|}. \quad (47)$$

Proof: On the one hand we have,

$$\begin{aligned} D_c f(x) = D_{b \oplus a} f(x) &= \left(\frac{1 - |b \oplus a|^2}{|1 - (b \oplus a)x|^2} \right)^{\frac{n-1}{2}} f(\varphi_{-b \oplus a}(x)) \\ &= \left(\frac{(1 - |a|^2)(1 - |b|^2)}{|1 - ab - (a + b)x|^2} \right)^{\frac{n-1}{2}} f(\varphi_{-b \oplus a}(x)). \end{aligned}$$

As $-\varphi_a(x) = \varphi_{-a}(-x)$ then $-b \oplus a = -\varphi_{-b}(a) = \varphi_b(-a) = (-b) \oplus (-a)$.

Moreover, by (2) we have that $\varphi_{-a} \circ \varphi_{-b}(x) = q \varphi_{(-b) \oplus (-a)}(x) \overline{q}$, with $q = \frac{1-ab}{|1-ab|}$, and thus

$$\varphi_{-b \oplus a}(x) = \overline{q}(\varphi_{-a}(\varphi_{-b}(x)))q. \quad (48)$$

Therefore, we have

$$\begin{aligned} D_b D_a R_q f(x) &= \left(\frac{1 - |b|^2}{|1 - bx|^2} \frac{1 - |a|^2}{|1 - a\varphi_{-b}(x)|^2} \right)^{\frac{n-1}{2}} f(\overline{q} \varphi_{-a}(\varphi_{-b}(x)) q) \\ &= \left(\frac{(1 - |a|^2)(1 - |b|^2)}{|1 - ab - (a + b)x|^2} \right)^{\frac{n-1}{2}} f(\varphi_{-b \oplus a}(x)), \text{ by (48).} \end{aligned}$$

■

By (47) we conclude that the operator $D_a R_q$ express the anisotropy character of the dilation.

Definition 5 *We define the anisotropy of a section $\sigma(te_n) = te_n \oplus g(t)e_{n-1}$ as*

$$\epsilon_g := \int_{-1}^1 \|D_{g(t)e_{n-1}} R_{q(t)} - I\| dt. \quad (49)$$

Since the operators $D_{g(t)e_{n-1}}$ and $R_{q(t)}$ are unitary we have an upper bound for our anisotropy, $0 \leq \epsilon_g \leq 4$. The concept of anisotropy has an important meaning, however, it is very difficult to compute the quantity ϵ_g . For that reason we propose an alternative definition using the generating function of the section $\sigma(te_n)$.

Definition 6 The p -deviation of the global continuous section $\sigma(te_n)$ is given by

$$\epsilon_{g,p}^* := \int_{-1}^1 \|\sigma(te_n) - te_n\|^p dt, \quad 1 \leq p < \infty$$

while we define the infinity deviation as $\epsilon_{g,\infty}^* := \sup_{t \in [-1,1[} \|\sigma(te_n) - te_n\|$.

Proposition 12 The p -deviation of $\sigma(te_n)$ is given by

$$\epsilon_{g,p}^* := \int_{-1}^1 \left(\frac{|g(t)|(1-t^2)}{\sqrt{1+t^2g(t)^2}} \right)^p dt, \quad 1 \leq p < \infty$$

and the infinity deviation is given by $\epsilon_{g,\infty}^* := \sup_{t \in [-1,1[} \frac{|g(t)|(1-t^2)}{\sqrt{1+t^2g(t)^2}}$.

For instance, for the sections $\sigma_\lambda(te_n)$ defined in Section 4 we obtain $\epsilon_{\lambda,p}^* := |\lambda|^p \int_{-1}^1 \left(\frac{1-t^2}{\sqrt{1+t^2\lambda^2}} \right)^p dt$, (e.g. $\epsilon_{\lambda,1}^* = \frac{|\lambda|(2\lambda^2 \ln(\lambda+\sqrt{1+\lambda^2}) - \lambda\sqrt{1+\lambda^2} + \ln(\lambda+\sqrt{1+\lambda^2}))}{\lambda^3}$) and $\epsilon_{\lambda,\infty}^* = |\lambda|$, while for the sections $\sigma_c(\phi)$ we obtain $\epsilon_{c,p}^* := |c|^p \int_0^\pi \sin^p \phi d\phi$ (e.g. $\epsilon_{c,1}^* = 2|c|$) and $\epsilon_{c,\infty}^* = |c|$.

The question of covariance of the spherical CWT (41) under rotations and dilations is very important. In the flat case, the usual n -dimensional CWT is fully covariant with respect to translations, rotations, and dilations, and this property is essential for applications, in particular, the covariance under translations. In fact, covariance is a general feature of all coherent state systems directly derived from a square integrable representation [4]. However, when the representation is only square integrable over a quotient of the group then no general theorem is available. In the case of conformlets the results are the following:

- The spherical CWT (41) is covariant under rotations on S^{n-1} : for any $s_1 \in \text{Spin}(n)$, the transform of the rotated signal $f(\overline{s_1}xs_1)$ is the function $W_\psi[f](\overline{s_1}s, \sigma(te_n))$.
- The spherical CWT (41) is not covariant under dilations. The wavelet transform of the dilated signal $D_{\sigma(t_1e_n)}f(x)$, is of the form $W_\psi[f](s, (\overline{s}\sigma(t_1e_n)s) \oplus (-\sigma(te_n)))$ as we can see by direct calculations.

In fact, if we consider the change of variables $\varphi_{-b}(x) = y$ ($\Leftrightarrow x = \varphi_b(y)$), with $dS_x = \left(\frac{1-|b|^2}{|1+by|^2} \right)^{\frac{n-1}{2}} dS_y$ we obtain

$$W_\psi[D_b f](s, a) = \int_{S^{n-1}} \left(\frac{1-|a|^2}{|1-a\overline{s}xs|^2} \right)^{\frac{n-1}{2}} \psi(\varphi_{-a}(\overline{s}xs)) \left(\frac{1-|b|^2}{|1-bx|^2} \right)^{\frac{n-1}{2}} f(\varphi_{-b}(x)) dS_x$$

$$\begin{aligned}
&= \int_{S^{n-1}} \left(\frac{(1-|a|^2)(1-|b|^2)^3}{|1-sa\bar{s}\varphi_b(y)|^2|1-b\varphi_b(y)|^2|1+by|^4} \right)^{\frac{n-1}{2}} \psi(\varphi_{-a}(\bar{s}\varphi_b(y)s))f(y)dS_y \\
&= \int_{S^{n-1}} \left(\frac{(1-|a|^2)(1-|b|^2)}{|1+by-sa\bar{s}(y-b)|^2} \right)^{\frac{n-1}{2}} \psi(\varphi_{(\bar{s}bs)\oplus(-a)}(\bar{s}ys))f(y)dS_y \\
&= W_\psi[f](s, (\bar{s}bs) \oplus (-a)), \quad \forall s \in \text{Spin}(n), \quad \forall a, b \in B^n.
\end{aligned}$$

As in general $(\bar{s}\sigma(t_1e_n)s) \oplus (-\sigma(te_n))$ is not an element of the section $\sigma(te_n)$, covariance under dilation does not hold. This is a consequence of the parameter space not being a group and is a general feature of coherent systems based on homogeneous spaces. For applications, of course, it is the covariance under rotations that is essential.

Acknowledgments

The author would like to thank Prof. Hans Feichtinger for his hospitality and support during a visit at NuHAG, Department of Mathematics, University of Vienna as well as Prof. J.-P. Antoine and Prof. P. Vandergheynst for the support and discussions during a stay at the *Institut de Physique Theorique* of *Université Catholique de Louvain*. The author is also very thankful to H. Rauhut, M. Fornasier, U. Kähler, P. Cerejeiras and N. Faustino for fruitful discussions. The work was supported by PhD-grant no. SFRH/BD/12744/2003 of *Fundação para a Ciência e a Tecnologia* of Portugal.

References

- [1] P. Abrial, Y. Moudden, M. Nguyen, J. L. Starck, *Curvelet transform on the sphere*, IEEE International Conference in Image Processing (ICIP 05), 2005.
- [2] P. Abrial, Y. Moudden, M. Nguyen, J. L. Starck, *Wavelets, Ridgelets and Curvelets on the sphere*, Astronomy and Astrophysics **446** (2006), 1191-1024.
- [3] L. Ahlfors, *Möbius transformations in $\mathbb{R}^n$ expressed through 2×2 matrices if Clifford numbers*, Complex Variables **5** (1986), 215-221.
- [4] S. T. Ali, J.-P. Antoine, J. P. Gazeau, *Coherent States, Wavelets and their Generalizations*, Springer-Verlag, Berlin, 2000.
- [5] S. T. Ali, J.-P. Antoine, J. P. Gazeau, *Square integrability of group representations on homogeneous spaces I. Reproducing triples and frames*, Ann. Inst. Henri Poincaré **55** (1991), 829-855.
- [6] S. T. Ali, J.-P. Antoine, J. P. Gazeau, *Square integrability of group representations on homogeneous spaces II. Coherent states and quasi-coherent states. The case of Poincaré group*, Ann. Inst. Henri Poincaré **55** (1991), 857-890.

- [7] J.-P. Antoine, L. Demanet, L. Jacques, P. Vandergheynst, *Wavelets on the sphere: Implementation and approximations*, Applied Comput. Harmon. Anal. **13** (2002), 177–200.
- [8] J.-P. Antoine, P. Vandergheynst, *Wavelets on the 2-sphere: A group-Theoretical Approach*, Appl. Comput. Harmon. Anal. **7** (1999), 262–291.
- [9] J.-P. Antoine, P. Vandergheynst, *Wavelets on the n -sphere and other manifolds*, J. Math. Physics **39** (1998), 3987–4008.
- [10] R. B. Barreiro, L. Cayón, E. M. González, J. L. Sanz, P. Vielva, *Detection of Non-Gaussianity in WMAP 1 Year Data*, Astrophysical Journal **609** (2004), 22–34.
- [11] L. Cayón, J. L. Sanz, E. M. González, A. J. Banday, F. Argüeso, J. E. Gallegos, K. M. Górski, G. Hinshaw *Spherical Mexican hat wavelet: an application to detect non-Gaussianity in the COBE-DMR maps*, Monthly Notices of the Royal Astronomical Society **326** (2001), 1243.
- [12] P. Cerejeiras, M. Ferreira, U. Kähler, F. Sommen, *Continuous wavelet transform and wavelet frames on the sphere using Clifford Analysis*, Comm. Pure Appl. Math., Vol. **6** no. 3 (2007), 619–641.
- [13] J. Cnops, *Hurwitz pairs and applications of Möbius transformations*, Habilitation dissertation, Universiteit Gent, Faculteit van de Wetenschappen, 1994.
- [14] S. Dahlke, M. Fornasier, H. Rauhut, G. Steidl, G. Teschke, *Generalized Coorbit Theory, Banach Frames and the Relation to α -Modulation Spaces*, Proc. London Math. Soc. (2005), to appear.
- [15] S. Dahlke, G. Steidl, G. Teschke, *Coorbit spaces and Banach Frames on Homogeneous Spaces with Applications to Analyzing Functions on Spheres*, Adv. Comput. Math. **21** no. 1–2 (2004), 147–180.
- [16] R. Delanghe, F. Sommen, V. Souček, *Clifford analysis and spinor-valued functions: a function theory for the Dirac operator*, Math. Appl. **53** (1990), Kluwer Acad. Publ., Dordrecht.
- [17] J. Fillmore, A. Springer, *Möbius groups over general fields using Clifford algebras associated with spheres*, Int. J. Theo. Phys. **29** (1990), 225–246.
- [18] W. Freeden, T. Gervens, M. Schreiner, *Constructive Approximation on the Sphere, with Applications to Geomathematics*, Clarendon Press, Oxford, 1997.
- [19] Y. Friedman, *Physical Applications of Homogeneous Balls*, Birkhäuser, 2005.
- [20] P. Frossard, I. Tosic, P. Vandergheynst *Progressive coding of 3-D objects based on overcomplete decompositions*, IEEE Trans. Circuits and Syst. Video Technol. **16**(11) (2006), 1338–1349.

- [21] S. Hanany, A. H. Jaffe, C. H. Lineweaver, L. Tenorio, *Applications of wavelets to the analysis of cosmic microwave background maps*, Monthly Notices of the Royal Astronomical Society **310** (1999), 823–834.
- [22] M. P. Hobson, A. N. Lasenby, J. D. McEwen, *A directional continuous wavelet transform on the sphere*, ArXiv, 2006, astro-ph/0609159.
- [23] M. P. Hobson, A. N. Lasenby, J. D. McEwen, D. J. Mortlock, *Fast directional continuous spherical transform algorithms*, IEEE Transactions on Signal Processing (2006), in press.
- [24] M. Holschneider, *Continuous wavelet transforms on the sphere*, J. Math. Phys. **37** (1996), 4156–4165.
- [25] L. Jacques, P. Vandergheynst, Y. Wiaux, *Correspondence principle between spherical and Euclidean wavelets*, Astrophysical Journal, **632** issue 1 (2005), 15–28.
- [26] M. K. Kinyon, O. Jones, *Loops and semidirect products*, Comm. Algebra **28**(9) (2000), 4137-4164.
- [27] V. Kisil, *Monogenic Calculus as an Intertwining Operator*, Bull. Belg. Math. Soc. Simon Stevin **11** no. 5 (2005), 739-757.
- [28] P. Schröder, W. Sweldens, *Spherical wavelets: Efficiently representing functions on the sphere*, Computer Graphics Proceedings (SIG- GRAPH 95) (1995), 161–172.
- [29] R. Takahashi, *Sur les représentations unitaires des groupes de Lorentz généralisés*, Bulletin de la Soc. Math. France **91** (1963), 289-433.
- [30] B. Torr sani, *Position-frequency analysis for signals defined on spheres*, Signal Proc. **43** (1995), 341–346.
- [31] A. A. Ungar, *Analytic Hyperbolic Geometry - Mathematical Foundations and Applications*, World Scientific, 2005.
- [32] A. A. Ungar, *Thomas precession and the parametrization of the Lorentz transformation group*, Found. Phys. Lett. **1** (1988), 57-89.
- [33] A. A. Ungar, *Thomas precession: its underlying gyrogroup axioms and their use in hyperbolic geometry and relativistic physics*, Found. Phys. **27**(6) (1997), 881-951.
- [34] N. Ja Vilenkin, A. U. Klimyk, *Representations of Lie Groups and Special Functions*, Kluwer Acad. Publisher, Vol. 2, 1993.